

# Compliance with expert population-based dietary guidelines and lower odds of carotid atherosclerosis in women: the Framingham Nutrition Studies<sup>1-3</sup>

Barbara E Millen, Paula A Quatromoni, Byung-Ho Nam, Michael J Pencina, Joseph F Polak, Ruth W Kimokoti, Jose M Ordovas, and Ralph B D'Agostino

## ABSTRACT

**Background:** Carotid stenosis, an indicator of subclinical atherosclerosis, predicts future coronary artery disease (CAD) and stroke and provides a noninvasive method to identify candidates for primary prevention. The relation between diet and stenosis is relatively unexplored, particularly in women.

**Objective:** We evaluated in women the association between nutrient intakes that were consistent with expert population-based dietary guidelines and carotid stenosis.

**Design:** We used prospective logistic regression analyses to evaluate relations between baseline nutrient intake and the presence of carotid stenosis at 4-y follow-up in 1123 women from the Framingham Offspring-Spouse study, after control for multiple CAD risk factors. We also developed multivariate models that were stratified by compliance with expert population-based dietary guidelines and smoking status.

**Results:** Baseline nutrient and risk factor profiles differed by women's compliance and smoking status. Dietary noncompliance and smoking were each associated with odds for stenosis that were 2.5-fold those of dietary compliance and nonsmoking. Odds were highest for dietary noncompliance in combination with smoking (odds ratio: 3.49; 95% CI: 1.67, 7.27).

**Conclusions:** Nutrient intake consistent with current expert population-based dietary guidelines and smoking abstinence are associated with lower odds of carotid atherosclerosis in women. Unique dietary and risk factor profiles of at-risk women suggest areas for targeted primary CAD prevention. *Am J Clin Nutr* 2005;82:174-80.

**KEY WORDS** Carotid stenosis, total and saturated fat, dietary cholesterol, smoking, preventive nutrition, coronary artery disease, CAD

## INTRODUCTION

Coronary artery disease (CAD) remains the leading cause of morbidity and mortality in adult Americans; it accounts for nearly 500 000 deaths annually and contributes to the \$300 billion in health care costs and lost individual productivity that is associated with cardiovascular disease (CVD) (1, 2). The primary prevention of CAD, particularly among men and women who have multiple CVD risk factors, is a national public health priority (3-7).

*Healthy People 2010* focuses on preventing CAD in addition to other chronic diseases (eg, stroke, obesity, diabetes, and cancer) and on promoting health by encouraging good nutrition and physical activity and the avoidance of smoking (4). The current US Department of Agriculture (USDA) *Dietary Guidelines for Americans 2000* (5) and the AHA [American Heart Association] *Dietary Guidelines* (6), which include lifestyle-related recommendations for CAD prevention, emphasize the population benefits of achieving and maintaining an overall healthy eating pattern, appropriate body weight, desirable lipid and blood pressure measurements, daily physical activity, and the avoidance of smoking. Specific population-based nutrient guidelines to lower CAD risk factor profiles recommend the following daily intakes:  $\leq 30\%$  of total energy as fat,  $< 10\%$  of energy as saturated fat, and  $< 300$  mg cholesterol (5, 6). Although these guidelines are similarly applicable to men and women, an even greater potential may exist for primary prevention in women because of the 10-15-y delay in the onset of CAD in women compared with men (7).

Over the past decade, increases in physician and public awareness of the benefits of reducing cholesterol concentrations, improvements in American dietary lipid and cholesterol intakes, and lower population serum cholesterol concentrations have contributed to declining rates of CAD. Nonetheless, recent reports (5, 7) emphasize that future success in reducing the national

<sup>1</sup> From the Department of Social and Behavioral Sciences, School of Public Health (BEM, PAQ, and RWK); the Department of Health Sciences, Sargent College of Health and Rehabilitation Sciences (PAQ); the Department of Socio-Medical Sciences (BEM), the Graduate Division of Medical Sciences, School of Medicine (BEM); and the Department of Mathematics (B-HN, MJP, and RBD), Boston University, Boston, MA; the Department of Radiology (JFP) and the Lipid Metabolism Laboratory, US Department of Agriculture Human Nutrition Research Center on Aging (JMO), Tufts University School of Medicine, Boston, MA; and the New England Medical Center, Boston, MA (JFP).

<sup>2</sup> Supported by grants no. R01-HL-60700 and N01-HC-38038 from the National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD.

<sup>3</sup> Address reprint requests to B Millen, Department of Social and Behavioral Sciences, Room 263W, Boston University School of Public Health, 715 Albany Street, Talbot 2 West, Boston, MA 02118. E-mail: [bmillen@bu.edu](mailto:bmillen@bu.edu). Received September 22, 2004.

Accepted for publication February 23, 2005.

burden of heart disease will depend on identifying effective prevention strategies, particularly those that focus on the following: primary prevention of CAD in the population, feasible targets for behavioral lifestyle modifications, and effective methods to promote long-term behavior change. Experts and clinicians also seek continuing evidence that the existing population-based lifestyle-related guidelines for CAD prevention promote CAD risk reduction (5, 7), including lowering the risk of preclinical CAD. Such research is particularly important in light of the dramatic rise in the proportion of American adults who are overweight and obese and the low rates of leisure-time physical activity noted in the US population (8, 9). In this study, we evaluated in women the relation between compliance with the USDA and AHA dietary guidelines and carotid artery stenosis, an indicator of subclinical atherosclerosis. Carotid stenosis predicts future fatal and nonfatal CAD and stroke (10–17) and offers a noninvasive method of identifying candidates for primary CAD-prevention activities (18, 19).

## SUBJECTS AND METHODS

### Subjects

The Framingham Study began in 1948 as a longitudinal, population-based study of CVD among residents of Framingham, MA (20; also RB D'Agostino, WB Kannel, unpublished observations, 1988–1989). In 1971, a second-generation cohort was recruited when some 5124 Framingham Study offspring and their spouses were invited to participate in the Framingham Offspring-Spouse (FOS) study (21).

Members of the FOS cohort are examined in the Framingham Study clinic, on average, every 4 y. They participate in a standardized protocol that involves a complete physical examination, laboratory tests, noninvasive diagnostic testing, and updating of medical histories and other pertinent information. At certain examinations, detailed dietary data are collected. The dietary and risk factor data reported here were collected among FOS women at Exam 5, between 1992 and 1996. Some 2007 women, aged 26–84 y, participated in this examination (76% of the original FOS cohort women). Women who were diagnosed with CVD at or before Exam 5 were excluded from these analyses (6.7%;  $n = 135$ ).

All participants provided written informed consent. This study was approved by the institutional review board at Boston University Medical Center.

### Nutrient intake and dietary compliance

Nutrient intake was estimated from 3-d dietary records collected with the use of a standardized, published method (22, 23). Participants were instructed by a registered dietitian in the clinic to record their intake over 2 weekdays and 1 weekend day, while adhering to their usual eating practices. Subjects were trained to estimate portion sizes with the use of a validated two-dimensional food portion visual aid (23). Some 68.5% of women ( $n = 1375$ ) completed the dietary record protocol. Dietary records were processed by trained coders who adhered to standardized protocols. Nutrient calculations were performed with the use of the MINNESOTA NUTRITION DATA SYSTEM software (version 2.6; Food Database 6A; Nutrient Database 23; Nutrition Coordinating Center, University of Minnesota, Minneapolis, MN; 24).

Subjects who were considered compliant had a 3-d dietary mean nutrient intake that met the population-based dietary guidelines for CAD risk reduction as set forth in the *AHA Dietary Guidelines* and that was consistent with the USDA's *Dietary Guidelines for Americans*, including a total of  $\leq 30\%$  of energy as fat (kcal),  $< 10\%$  of energy as saturated fat (kcal), and  $< 300$  mg dietary cholesterol (4–6). Noncompliant subjects consumed diets that failed to meet  $\geq 1$  of these guidelines. We compared these dietary guidelines with the medical nutrition therapy recommendations (7), to assess the potential benefits associated with population-based recommendations in relation to an early indicator of atherosclerotic heart disease.

### Assessment of carotid atherosclerosis

Among 1283 women who provided complete dietary records and who were free of CVD at baseline, the presence of carotid atherosclerosis was assessed at follow-up at Exam 6 (1996–1999). Carotid ultrasound scanning studies were obtained for 1137 subjects (89% of the total cohort). Missing measurements were exclusively due to logistic constraints at the clinic (eg, unavailability of the ultrasound scanning device or the sonographer during the scheduled visit). Reliability studies were ongoing during data acquisition.

Ultrasound imaging studies were conducted with a high-resolution linear-array 5.0 MHz transducer and color Doppler ultrasound scanning device (Toshiba SSH-140; Toshiba Medical Systems, Tustin, CA). Imaging was performed with the subject's head rotated 45 degrees away from the side being studied, according to a standard protocol. Two gray-scale images were taken at the level of the common carotid artery bulb, and 2 additional images were obtained in the proximal 2 cm of the internal carotid artery. One image of the respective image pairs was acquired with the probe held at 45 degrees from horizontal. For the second image, the sonographer was instructed to position the transducer to best identify any focal lesions. All images were gated to the R-wave of an electrocardiogram, and both sides of the neck were imaged. Images were directly transferred into a computer workstation through a frame-grabber board.

Color Doppler imaging and pulsed Doppler waveforms were used to evaluate blood flow velocities in the proximal internal carotid arteries. Angle-corrected Doppler velocity waveforms were acquired in the proximal internal carotid artery at the site of highest velocity as identified on a color Doppler image. Peak systolic velocities were measured from these tracings. A certified reader reviewed the acquired digital images and made a subjective estimate of the degree of internal carotid artery narrowing, which was graded as 0%, 1–24%, or 25–49% when Doppler-derived peak systolic velocities in the internal carotid artery were  $< 150$  cm/s. Internal carotid artery disease was characterized by the maximum stenosis observed on the right or left side and was categorized as 0% (no lesions), 1–24%, 25–49% (focal lesions causing  $< 50\%$  diameter stenosis), or  $\geq 50\%$  (lesions causing  $\geq 50\%$  diameter stenosis). Absence of blood flow corresponded to a total occlusion.

### CVD risk factor measurements

CVD risk factors are routinely measured at all Framingham Study examinations (25). All lipid analyses were performed at the Framingham Study laboratory, which participates in the Standardization Program of the Centers for Disease Control and



Prevention and the National Heart, Lung, and Blood Institute Lipid Research Clinics. Venous blood was drawn from all subjects after a 12–24-h fast. Total and HDL-cholesterol concentrations were measured by automated enzymatic methods (26, 27). The cholesterol content of LDL cholesterol was estimated by the method of Friedewald et al (28). Triacylglycerol concentrations were measured enzymatically (26). Blood pressure was determined by duplicate measurements of the subject's left arm made with the use of a mercury sphygmomanometer while the subject was in a sitting position. Body mass index (BMI; in kg/m<sup>2</sup>) was calculated from height and weight values measured in the clinic. Diabetes was defined as a history of use of insulin or an oral hypoglycemic agent or a fasting blood glucose concentration of  $\geq 7.8$  mmol/L ( $\geq 140$  mg/dL).

Physical activity was measured with the use of a standardized questionnaire to determine estimates of activity in a usual day, based on a 24-h history. A physical activity index was calculated from the number of hours spent doing specific activities that were categorized (ie, sedentary, slight, moderate, or heavy) and weighted according to the oxygen consumption required to perform them (29). Physical activity index scores ranged from 24 (total bed rest) to 120. Because physical activity was not measured at FOS Exam 3, Exam 2 values were substituted in these analyses, according to the analytic approach routinely used in the Framingham study. Menopausal status and cigarette smoking was self-reported. Pack-years of cigarette smoking were defined as the number of packs of cigarettes smoked per day multiplied by the total number of years a person smoked. This variable takes into account the duration of smoking as well as the amount of cigarette consumption.

### Statistical analysis

Our primary objective was to determine whether compliance with the current population-based dietary guidelines for CAD risk reduction (as defined earlier) was related to the presence of carotid atherosclerosis at follow-up. The endpoint of interest was carotid artery stenosis, defined by focal lesions of  $\geq 25\%$  in either the right or left internal carotid artery, in accordance with our previously used threshold (30–32). Previous research suggested potential interaction between diet and smoking (32, 33); thus, we decided to test for such an interaction. Because the null hypothesis (ie, no interaction) could not be rejected ( $P = 0.10$ ), we decided to stratify by compliance and smoking status and created 4 subgroups: compliant nonsmokers, noncompliant nonsmokers, compliant smokers, and noncompliant smokers.

For descriptive purposes, age-adjusted mean baseline Exam 5 CVD risk factors and nutrient intake profiles were computed for the 4 subgroups. Analysis of covariance was used for calculating the least-squares means of continuous variables by using PROC GLM in SAS (34). For categorical variables, age-adjusted proportions were computed by using the age-adjusted logistic regression (PROC LOGISTIC) (34). If the interaction between smoking and compliance was significant, we compared the 4 subgroups by using Bonferroni's correction for multiple testing (34). In multivariate analyses, we examined the relation between dietary compliance and carotid atherosclerosis at follow-up with dietary compliance and smoking subgroups by using compliant nonsmokers as the reference group. Odds ratios (ORs) were calculated by using logistic regression in which CVD risk factors were considered separately in age- and energy-adjusted models. Models considered a range of CVD risk factors, including age,

systolic and diastolic blood pressures, total cholesterol, LDL cholesterol, the ratio of total to HDL cholesterol, plasma triacylglycerols, BMI, physical activity level, menopausal status, and conditions that included obesity, diabetes, and hypertension. In addition, education level (high school or below compared with more than high school) was considered as a proxy indicator of socioeconomic status. The final multivariate model reported here is limited to those variables that were identified as important predictors or potential confounders of the relation of interest. The sample used in the analyses consisted of women with no missing values for carotid stenosis, dietary compliance, smoking, and the covariates retained in the final multivariate model ( $n = 1123$ ). Analyses were carried out by use of SAS software (version 8.2; SAS Institute Inc, Cary, NC).

### RESULTS

Among 1123 women studied, 360 (32%, both smokers and nonsmokers) had baseline age-adjusted mean daily nutrient intakes that complied with all 3 criteria of the USDA and AHA dietary guidelines for CAD risk reduction (5, 6): a total of  $\leq 30\%$  of energy as fat,  $< 10\%$  of energy as saturated fat, and  $< 300$  mg dietary cholesterol (**Table 1**). Some 68% of FOS women ( $n = 763$ ) had nutrient profiles that failed to meet  $\geq 1$  of these current population dietary recommendations (noncompliant smokers and nonsmokers). Compliant and noncompliant subjects (both smokers and nonsmokers) also had markedly different intakes of a wider range of nutrients and CVD risk factors. Specifically, the diets of noncompliant women were higher in total energy, dietary lipids, cholesterol, and sodium; less concentrated in carbohydrate and fiber; and less nutrient-dense overall than were the diets of women who were compliant. Smokers (both compliant and noncompliant) had higher alcohol intakes than did nonsmokers. Noncompliant smokers had the lowest intakes of folate.

The baseline CVD characteristics of FOS women are presented in **Table 2**. Noncompliant women were younger and had lower total:HDL cholesterol than did compliant women. Compared with compliant smokers, noncompliant smokers had higher levels of cigarette exposure at baseline. The means of systolic and diastolic blood pressure; BMI; total, HDL, and LDL cholesterol; physical activity; and the rates of diabetes did not differ significantly between compliant and noncompliant subjects.

The multivariate models, stratified by the combination of dietary compliance and smoking exposure, are presented in **Table 3**. Compliant women who never smoked were the reference group. Compared with nonsmoking subjects who complied with the current dietary guidelines, noncompliant women who never smoked had 2.5-fold higher odds of carotid stenosis. The odds of carotid stenosis in women with compliant diets at baseline but who smoked was similar to the odds in noncompliant women who smoked. Noncompliant women who also smoked at baseline had odds of carotid atherosclerosis at 4 y of follow-up 3.5 times those in the reference group.

### DISCUSSION

This is the first cohort study in women to examine the relation between compliance with population-based dietary guidelines for CAD prevention and the incidence of carotid stenosis, a subclinical marker of systemic atherosclerosis (10, 11, 35, 36)

TABLE 1

Subject nutrient intake by dietary compliance and smoking status at baseline (1992–1996)<sup>1</sup>

Nutrient intake	Compliant nonsmokers <sup>2</sup> (n = 157)	Compliant smokers <sup>2</sup> (n = 203)	Noncompliant nonsmokers <sup>3</sup> (n = 311)	Noncompliant smokers <sup>3</sup> (n = 452)	P <sup>4</sup>	P <sup>5</sup>	P <sup>6</sup>
Energy (kcal)	1510.6 ± 38.4 <sup>7</sup>	1517.1 ± 33.8	1700.1 ± 27.3	1694.5 ± 22.7		0.96	<0.0001
Total fat (% of energy)	24.1 ± 0.39	23.7 ± 0.34	35.6 ± 0.28	35.7 ± 0.23		0.83	<0.0001
Saturated fat (% of energy)	7.3 ± 0.19	7.1 ± 0.17	12.1 ± 0.13	12.0 ± 0.11		0.59	<0.0001
Polyunsaturated fat (% of energy)	5.6 ± 0.18	5.7 ± 0.16	7.6 ± 0.13	7.6 ± 0.11		0.81	<0.0001
Monounsaturated fat (% of energy)	8.9 ± 0.18	8.6 ± 0.16	13.2 ± 0.13	13.3 ± 0.11		0.98	<0.0001
Carbohydrate (% of energy)	57.7 ± 0.58	56.4 ± 0.51	47.4 ± 0.42	45.8 ± 0.35		0.0005	<0.0001
Protein (% of energy)	18.4 ± 0.31	19.1 ± 0.27	16.9 ± 0.22	17.2 ± 0.18		0.081	<0.0001
Alcohol (g)	4.2 ± 0.82	7.1 ± 0.72	4.1 ± 0.58	6.7 ± 0.49		<.0001	0.67
Cholesterol (mg)	143.1 ± 8.1	150.1 ± 7.1	228.5 ± 5.7	231.8 ± 4.8		0.46	<0.0001
Dietary fiber (g)	17.5 ± 0.54	17.9 ± 0.47	15.2 ± 0.38	14.5 ± 0.32		0.35	<0.0001
Sodium (mg)	2304.6 ± 77.4	2418.9 ± 68.0	2715.7 ± 55.0	2802.0 ± 45.7		0.105	<0.0001
Folacin (μg)	289.3 ± 10.6 <sup>a</sup>	302.0 ± 9.3 <sup>a</sup>	263.2 ± 7.5 <sup>b</sup>	240.2 ± 6.2 <sup>c</sup>	0.037		
Vitamin C (mg)	136.6 ± 5.3	129.5 ± 4.7	99.4 ± 3.8	97.0 ± 3.1		0.32	<0.0001
Vitamin E (mg)	8.1 ± 0.40	8.0 ± 0.35	9.4 ± 0.28	8.9 ± 0.24		0.28	0.0011
Vitamin B-6 (mg)	1.8 ± 0.06 <sup>a</sup>	1.9 ± 0.05 <sup>a</sup>	1.7 ± 0.04 <sup>b</sup>	1.6 ± 0.03 <sup>b</sup>	0.068		
β-Carotene (μg)	4774.2 ± 353.3	4813.4 ± 310.5	4085.7 ± 250.8	3734.3 ± 208.7		0.40	0.0013
Calcium (mg)	672.8 ± 25.6	658.4 ± 22.5	688.7 ± 18.2	658.8 ± 15.1		0.20	0.73
Potassium (mg)	2743.4 ± 67.9	2747.8 ± 59.6	2552.9 ± 48.2	2533.1 ± 40.1		0.82	0.0002
Magnesium (mg)	276.5 ± 7.5	284.5 ± 6.6	265.6 ± 5.3	259.2 ± 4.4		0.76	0.0016

<sup>1</sup> The sample includes 1123 women free of cardiovascular disease at baseline with complete data on covariates. Analysis of covariance was used to obtain the age-adjusted means for continuous variables and to identify significant differences. Logistic regression was used to obtain the age-adjusted proportions and to identify subgroups that differ significantly. Means in a row with different superscript letters are significantly different,  $P < 0.05$  (Bonferroni's corrected  $t$  tests).

<sup>2</sup> Three-day mean intake met the population-based dietary guidelines from the US Department of Agriculture (USDA) and the American Heart Association (AHA) for coronary artery disease risk reduction for all 3 of the following nutrients: total fat  $\leq 30\%$  of energy (kcal), saturated fat  $< 10\%$  of energy (kcal), and cholesterol  $< 300$  mg.

<sup>3</sup> Three-day mean intake failed to meet the population-based USDA and AHA dietary guidelines for coronary artery disease risk reduction for  $\geq 1$  of the 3 nutrients of interest.

<sup>4</sup> The smoking  $\times$  noncompliance interaction.

<sup>5</sup> The main effect of smoking if the smoking  $\times$  noncompliance interaction was not significant.

<sup>6</sup> The main effect of noncompliance if the smoking  $\times$  noncompliance interaction was not significant.

<sup>7</sup>  $\bar{x} \pm SE$  (all such values).

that predicts future CAD and cerebrovascular events (10, 12–17, 37, 38). More than two-thirds of the women in the FOS Study had nutrient intakes that failed to meet the current population-based nutritional guidelines (from USDA and AHA) at baseline in 1992–1996. Dietary noncompliance and cigarette smoking each more than doubled the odds for stenosis seen with dietary compliance. Noncompliant women who smoked experienced a tripling of the odds of carotid stenosis.

These findings are consistent with emerging evidence on lifestyle behaviors in relation to carotid stenosis, CAD risk, and other health outcomes in adult populations. We have reported on the associations between the 5 distinct dietary patterns of FOS Study women and the presence of carotid stenosis at 12 y of follow-up (32). After multivariate adjustment for age, systolic blood pressure, BMI, total:HDL cholesterol, and pack-years of smoking, women whose dietary patterns were characterized as “empty calorie” (ie, more concentrated in high-fat, high-sugar foods and lower in fruit and vegetables and micronutrient-rich foods) were compared with women with more heart-healthy diets (ie, higher intakes of vegetables, fruit, and low-fat foods), and it was found that the former group had a significantly higher risk of carotid stenosis at 12-y follow-up [relative risk (RR): 2.28; 95% CI: 1.12, 4.62]. In a follow-up study, stratified by smoking status, women in the heart-healthy cluster who had never smoked had a

significantly lower risk of carotid stenosis (OR: 0.17; 95% CI: 0.07, 0.36) than did women in the less heart-healthy group at 12-y follow-up (39).

McCullough et al (40) used the 10-item Healthy Eating Index (HEI) (41) to assess the relation between compliance with the USDA's *Dietary Guidelines for Americans* (42) and the *Food Guide Pyramid* (43) and various chronic disease endpoints. Women in the highest quintile of HEI score (which indicated a more desirable eating pattern) had 14% lower CVD risk over 12 y than did women in the lowest HEI quintile. Higher HEI was not associated with lower cancer risk or overall chronic disease risk in these women.

Stampfer et al (44) examined healthy lifestyle behaviors in relation to the development of coronary events (death due to CAD or nonfatal myocardial infarction) in women. Combinations of lifestyle behaviors were based on the following risk factors: diet (low in *trans* fat and glycemic load; high in cereal fiber, marine n-3 fatty acids, and folate; and a high ratio of polyunsaturated to saturated fat), smoking abstinence, physical activity ( $\geq 30$  min/d of moderate to vigorous exercise), BMI  $< 25$ , and alcohol consumption  $\geq 5$  g/d. At 14-y follow-up, women with healthier diets (upper 2 quintiles) who did not smoke and who exercised had a reduction of almost 60% in coronary events (RR: 0.43; 95% CI: 0.33, 0.55). The addition of

**TABLE 2**Subject cardiovascular disease (CVD) risk factor characteristics by dietary compliance and smoking status at baseline (1992–1996)<sup>1</sup>

CVD risk factor characteristics	Compliant nonsmokers <sup>2</sup> (n = 157)	Compliant smokers <sup>2</sup> (n = 203)	Noncompliant nonsmokers <sup>3</sup> (n = 311)	Noncompliant smokers <sup>3</sup> (n = 452)	P <sup>4</sup>	P <sup>5</sup>	P <sup>6</sup>
Age (y)	56.8 ± 9.1 <sup>7</sup>	56.2 ± 9.1	56.1 ± 10.6	54.0 ± 9.2		0.0071	0.011
Systolic blood pressure (mm Hg)	124.9 ± 1.4	125.5 ± 1.2	123.5 ± 1.0	123.0 ± 0.8		0.90	0.060
Diastolic blood pressure (mm Hg)	73.0 ± 0.78	72.4 ± 0.68	72.4 ± 0.55	72.2 ± 0.46		0.54	0.56
BMI (kg/m <sup>2</sup> )	26.4 ± 0.42	25.7 ± 0.37	26.6 ± 0.29	26.7 ± 0.25		0.47	0.056
Total cholesterol (mg/dL)	207.9 ± 2.7	209.8 ± 2.4	203.3 ± 1.9	208.5 ± 1.6		0.043	0.21
HDL cholesterol (mg/dL)	55.3 ± 1.2	56.0 ± 1.0	57.3 ± 0.85	57.3 ± 0.70		0.81	0.090
Total:HDL cholesterol	4.0 ± 0.10	4.1 ± 0.09	3.7 ± 0.07	3.9 ± 0.06		0.099	0.0088
LDL cholesterol (mg/dL)	125.4 ± 2.5	125.7 ± 2.2	122.5 ± 1.8	125.0 ± 1.5		0.36	0.44
Diabetes (%)	0.05 ± 0.02	0.05 ± 0.01	0.04 ± 0.01	0.04 ± 0.01		0.99	0.35
Cigarette exposure (pack-years)	0.0 <sup>a</sup>	14.6 ± 1.16 <sup>b</sup>	0.0 <sup>a</sup>	21.1 ± 0.78 <sup>c</sup>	0.0025		
Physical activity index	33.7 ± 0.40	33.8 ± 0.35	33.7 ± 0.28	33.6 ± 0.23		0.82	0.73

<sup>1</sup> The sample includes 1123 women free of CVD at baseline with complete data on covariates. Analysis of covariance was used to obtain the age-adjusted means for continuous variables and to identify significant differences. Logistic regression was used to obtain the age-adjusted proportions and to identify subgroups that differ significantly. Means in a row with different superscript letters are significantly different,  $P < 0.05$  (Bonferroni's corrected  $t$  tests).

<sup>2</sup> Three-day mean intake met the population-based dietary guidelines from the US Department of Agriculture (USDA) and American Heart Association (AHA) for coronary artery disease (CAD) risk reduction for all 3 of the following nutrients: total fat  $\leq 30\%$  of energy (kcal), saturated fat  $< 10\%$  of energy (kcal), and cholesterol  $< 300$  mg.

<sup>3</sup> Three-day mean intake failed to meet the population-based USDA and AHA dietary guidelines for CAD risk reduction for  $\geq 1$  of the 3 nutrients of interest.

<sup>4</sup> The smoking  $\times$  noncompliance interaction.

<sup>5</sup> The main effect of smoking if the smoking  $\times$  noncompliance interaction was not significant.

<sup>6</sup> The main effect of noncompliance if the smoking  $\times$  noncompliance interaction was not significant.

<sup>7</sup>  $\bar{x} \pm SE$  (all such values).

BMI  $< 25$  or both favorable BMI and alcohol intakes to this constellation of risk factors lowered women's risk of coronary events further (RR: 0.34; 95% CI: 0.23, 0.52 and RR: 0.17; 95% CI: 0.07, 0.41, respectively).

In the current study, FOS Study women who had carotid stenosis at 4-y follow-up had distinct patterns of nutrient intake, including higher total and saturated fat and sodium intakes and lower consumption of fiber, folacin, vitamins C and B-6, potassium, and magnesium. They also had a higher burden of CVD risk factors, including elevated systolic blood pressure, dyslipidemia, diabetes, and smoking exposure. Physical activity levels were similar in women with and without stenosis but were low, on average, in this cohort. Consistent with these data, Kuller et al (45) showed that the presence of subclinical atherosclerosis in women was associated with modifiable CVD risk factors, including cigarette smoking, higher LDL- and lower HDL-cholesterol concentrations, and higher systolic blood pressure measurement and blood glucose concentrations. From the standpoint of primary CAD prevention, the behavioral and CVD risk

factor profiles characteristic of women at risk of subclinical disease suggest certain targets for population-based interventions. They also provide a rationale for more aggressive intervention among those persons at high risk of subclinical disease (45).

The potential complexity of behavioral intervention to promote CAD risk reduction is underscored by these findings. Indeed, about two-thirds of women in the FOS Study population (ie, the diet-noncompliant smokers and nonsmokers) were unable to follow the current population-based dietary guidelines for CVD risk reduction and CAD prevention. The distinct nutrient intake profile of noncompliant women (dietary fat- and sodium-rich and poor micronutrient density) in relation to women with compliant diets points to the potential importance of targeting prevention messages and interventions to the specific lifestyle patterns of women.

Our data suggest that dietary noncompliance and cigarette smoking conferred a similar, 2.5-fold greater risk of stenosis that was independent of other CVD risk factors. Furthermore, the

**TABLE 3**Odds ratios (ORs) for stenosis in women stratified by dietary compliance and smoking status (1996–1999)<sup>1</sup>

	Coefficient estimate	P <sup>2</sup>	OR (95% CI) <sup>3</sup>
Compliant nonsmokers (n = 157)	Reference group		1.00
Noncompliant nonsmokers (n = 311)	0.9298	0.0171	2.53 (1.18,5.44)
Compliant smokers (n = 203)	0.9276	0.0214	2.53 (1.15,5.57)
Noncompliant smokers (n = 452)	1.2491	0.0009	3.49 (1.67,7.27)

<sup>1</sup> The sample includes 1123 women free of cardiovascular disease at baseline with complete data on covariates. Dietary compliance is defined as 3-d mean intake that meets the population-based dietary guidelines from the US Department of Agriculture and the American Heart Association for coronary artery disease risk reduction for all 3 of the following nutrients: total fat  $\leq 30\%$  of energy (kcal), saturated fat  $< 10\%$  of energy (kcal), and cholesterol  $< 300$  mg. Women who were noncompliant had 3-d mean intakes that failed to meet the guidelines for  $\geq 1$  of the 3 nutrients of interest.

<sup>2</sup> A formal test for the effect of the smoking status  $\times$  dietary compliance interaction on stenosis was conducted and yielded  $P = 0.10$ .

<sup>3</sup> Multivariate logistic regression model adjusted for age, energy intake, systolic blood pressure, total:HDL cholesterol, BMI, and diabetes.

combination of poor eating behavior and smoking appears to be associated with the highest risk of carotid atherosclerosis. Combined lifestyle risks are increasingly recognized (32, 44) but have yet to be formally integrated on a widespread basis into targeted, combined risk-reduction interventions. The high rates of smoking among women with unfavorable dietary patterns also should be recognized and targeted as a potential barrier to changes in nutrition behavior.

The assessment of carotid stenosis by high-resolution B-mode ultrasonography is accurate in diagnosing subclinical CVD (18, 19, 35, 45) and provides a noninvasive method of identifying potential candidates for primary prevention activities. It may also offer an approach for evaluating interventions that are aimed at delaying the development of atherosclerosis (10, 45).

Our observations were established in a cohort of women, most of whom are white residents of a western Boston suburban community. Although this population may somewhat limit the generalizability of our findings, we note that the Framingham Study models for CVD risk have been repeated and confirmed in domestic and international populations (46–49). Our data encourage further research on the relations in women among dietary patterns, subclinical CVD, and other health outcomes. 

BEM provided overall direction to this research, including the identification of research questions and hypotheses, review of data analyses, and the development of all aspects of the manuscript, which included the description of the conceptual framework and context of the research, the presentation of results, and the interpretation and discussion of the findings. PAQ was involved in all aspects of this research and manuscript preparation, particularly the presentation of research methods, the results, and their interpretation and the discussion of findings within the context of epidemiologic literature. B-HN and MJP performed all the statistical analyses related to this research and prepared the analytic methods section of this manuscript. JFP was responsible for providing clinical expertise on the use of the carotid stenosis data sets in these analyses, for the presentation of the clinical methods and findings, and for the discussion and interpretation of the stenosis results in this manuscript. RWK was responsible for the review and the development of all aspects of the manuscript in preparation for publication. JMO, as principal investigator of one of the funded research projects that supported this work, provided significant consultation and advice to the research team during the preparation of research for publication. RBD, as senior biostatistician and head of the Statistical Consulting Unit at the Framingham Study, provided oversight on all aspects of the statistical analyses, the analytic components of this manuscript, and the interpretation of the data. None of the authors had a personal or financial conflict of interest.

## REFERENCES

- American Heart Association. Heart and stroke statistics—2004 update. Internet: <http://www.americanheart.org> (accessed 1 August 2004).
- National Heart, Lung, and Blood Institute. Fact book fiscal year 2001. Bethesda, MD: National Heart, Lung, and Blood Institute, National Institutes of Health, 2001.
- National Center for Health Statistics. *Healthy People 2000* final review. Hyattsville, MD: Public Health Service, 2001.
- US Department of Health and Human Services. *Healthy people 2010: understanding and improving health*. 2nd ed. Washington, DC: US Government Printing Office, 2000.
- Dietary guidelines for Americans 2000. 5th ed. Center for Nutrition Policy and Promotion. Internet: <http://www.usda.gov/cnpp> (accessed 1 August 2004).
- Krauss RM, Eckel RH, Howard B, et al. AHA dietary guidelines: revision 2000. A statement for healthcare professionals from the Nutrition Committee of the American Heart Association. *Circulation* 2000;102:2284–99.
- NCEP Expert Panel. Executive summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001;285:2486–97.
- US Department of Health and Human Services. Physical activity and health: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 1996.
- Centers for Disease Control and Prevention, US Department of Health and Human Services. CDC fact book 2000/2001. Atlanta, GA: CDC, 2000.
- Salonen JT, Salonen R. Ultrasonographically assessed carotid morphology and the risk of coronary heart disease. *Arterioscler Thromb* 1991;11:1245–9.
- Selhub J, Jacques PF, Bostom AG, et al. Association between plasma homocysteine concentrations and extracranial carotid-artery stenosis. *N Engl J Med* 1995;332:286–91.
- Kuller LH, Shemanski L, Psaty BM, et al. Subclinical disease as an independent risk factor for cardiovascular disease. *Circulation* 1995;92:720–6.
- Ogren M, Hedblad B, Isacson S, et al. Ten year cerebrovascular morbidity and mortality in 68 year old men with asymptomatic carotid stenosis. *BMJ* 1995;310:1294–8.
- Longstreth WT Jr, Shemanski L, Lefkowitz D, et al. Asymptomatic internal carotid artery stenosis defined by ultrasound and the risk of subsequent stroke in the elderly: the Cardiovascular Health Study. *Stroke* 1998;29:2371–6.
- Inzitari D, Eliasziw M, Gates P, et al. The causes and risk of stroke in patents with asymptomatic internal-carotid-artery stenosis. *N Engl J Med* 2000;342:1693–700.
- Molloy J, Markus HS. Asymptomatic embolization predicts stroke and TIA risk in patients with carotid artery stenosis. *Stroke* 1999;30:1440–3.
- European Carotid Surgery Trialists' Collaborative Group. Risk of stroke in the distribution of an asymptomatic carotid artery. *Lancet* 1995;345:209–12.
- AbuRahma AF, Pollack JA, Robinson PA, et al. The reliability of color duplex ultrasound in diagnosing total carotid artery occlusion. *Am J Surg* 1997;174:185–7.
- Modaresi KB, Cox TCS, Summers PE, et al. Comparison of intra-arterial digital subtraction angiography, magnetic resonance angiography and duplex ultrasonography for measuring carotid artery stenosis. *Br J Surg* 1999;86:1422–6.
- Dawber TR. The Framingham Study. The epidemiology of atherosclerotic disease. Cambridge, MA: Harvard University Press, 1980.
- Kannel WB, Feinleib M, McNamara PM, et al. An investigation of coronary heart disease in families: the Framingham Offspring Study. *Am J Epidemiol* 1979;110:281–90.
- Posner BM, Martin-Munley SS, Smigelski C, et al. Comparison of techniques for estimating nutrient intake: the Framingham Study. *Epidemiology* 1992;3:171–7.
- Posner BM, Smigelski C, Duggal A, Morgan JL, Cobb J, Cupples LA. Validation of two-dimensional models for estimation of portion size in nutrition research. *J Am Diet Assoc* 1992;92:738–41.
- Schakel SF, Sievert YA, Buzzard IM. Sources of data for developing and maintaining a nutrient database. *J Am Diet Assoc* 1988;88:1268–71.
- Cupples LA, D'Agostino RB. Some risk factors related to the annual incidence of cardiovascular disease and death using pooled repeated biennial measurements: Framingham Heart Study, 30-year follow-up. In: Kannel WB, Wolf PA, Garrison RJ, eds. *The Framingham Study, an epidemiological investigation of cardiovascular disease*. Washington, DC: Department of Health and Human Services, 1987. (NIH publication 87-2703; NTIS PB87-177499.)
- McNamara JR, Schaefer EJ. Automated enzymatic standardized lipid analyses for plasma and lipoprotein fractions. *Clin Chem Acta* 1987;166:1–8.
- Warnick GR, Benderson J, Albers JJ. Dextran sulfate-magnesium precipitation procedure for quantification of high-density lipoprotein cholesterol. *Clin Chem* 1982;28:1379–82.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without the use of preparative ultracentrifuge. *Clin Chem* 1972;18:499–502.
- Kannel WB, Sorlie P. Some health benefits of physical activity. *The Framingham Study*. *Arch Intern Med* 1979;139:857–61.

30. Wilson PWF, D'Agostino RB, Levy D, et al. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998;97:1837-47.
31. Selhub J, Jacques PF, Bostom AG, et al. Relationship between plasma homocysteine, vitamin status and extracranial carotid-artery stenosis in the Framingham Study population. *J Nutr* 1996;126:1258S-65S.
32. Millen BE, Quatromoni PA, Nam B-H, O'Horo CE, Polak JF, D'Agostino RB. Dietary patterns and the odds of carotid atherosclerosis in women. The Framingham Nutrition Studies. *Prev Med* 2002;35:540-7.
33. Millen BE, Quatromoni PA, Copenhafer DL, Demissie S, O'Horo CE, D'Agostino RB. Validation of a dietary pattern approach for evaluating nutritional risk. The Framingham Nutrition Studies. *J Am Diet Assoc* 2001;101:187-94.
34. SAS Institute, Inc. SAS user's guide. Version 6, vol 1 and vol 2. 4th ed. Cary, NC: SAS Institute, 1989.
35. Wilson PW, Hoeg JM, D'Agostino RB, et al. Cumulative effects of high cholesterol levels, high blood pressure, and cigarette smoking on carotid stenosis. *N Engl J Med* 1997;377:516-22.
36. Salonen JT. Is there a continuing need for longitudinal epidemiological research? The Kuopio Ischaemic Heart Disease Risk Factor Study. *Ann Clin Res* 1988;20:46-50.
37. Autret A, Saudeau D, Bertrands PH, Pourcelot L, Marchal C, de Boissvilliers S. Stroke risk in patients with carotid stenosis. *Lancet* 1987; 1:888-90.
38. Mansour M, Littooy FN, Watson WC, et al. Outcome of moderate carotid artery stenosis in patients who are asymptomatic. *J Vasc Surg* 1999;29:217-27.
39. Millen BE, Quatromoni PA, Nam BH, et al. Dietary patterns, smoking, and subclinical heart disease in women: opportunities for primary prevention from the Framingham Nutrition Studies. *J Am Diet Assoc* 2004; 104:208-14.
40. McCullough ML, Feskanich D, Stampfer MJ, et al. Adherence to the Dietary Guidelines for Americans and risk of major chronic disease in women. *Am J Clin Nutr* 2000;72:1214-22.
41. Kennedy ET, Ohls J, Carlson S, Fleming K. The Healthy Eating Index: design and applications. *J Am Diet Assoc* 1995;95:1103-8.
42. US Department of Agriculture, US Department of Health and Human Services. Nutrition and your health: dietary guidelines for Americans. Washington, DC: US Government Printing Office, 2000.
43. US Department of Agriculture. The food guide pyramid. Hyattsville, MD: Human Nutrition Information Service, August 1992. (Home and Garden Bulletin no. 252.)
44. Stampfer MJ, Hu FB, Manson JE, et al. Primary prevention of coronary heart disease in women through diet and lifestyle. *N Engl J Med* 2000; 343:16-22.
45. Kuller L, Borhani N, Furberg C, et al. Prevalence of subclinical atherosclerosis and cardiovascular disease and association with risk factors in the Cardiovascular Health Study. *Am J Epidemiol* 1994;139:1164-79.
46. Gordon T, Garcia-Palmieri MR, Kagan A, Kannel WB, Schiffman J. Differences in coronary heart disease in Framingham, Honolulu and Puerto Rico. *J Chronic Dis* 1974;27:329-44.
47. McGee D, Gordon T. The Framingham Study applied to four other U.S.-based epidemiological studies of cardiovascular disease (section no. 31). Bethesda, MD: US Department of Health, Education, and Welfare, 1976. (NIH publication 76-1083.)
48. Brand RJ, Rosenman RH, Scholtz RI. Multivariate prediction of coronary heart disease in the Western Collaborative Group Study compared to the findings of the Framingham Study. *Circulation* 1976;53:348-55.
49. Leaverton PE, Sorlie PD, Kleinman JC, et al. Representativeness of the Framingham risk model for coronary heart disease mortality; a comparison with a national cohort study. *J Chronic Dis* 1987;40:775-84.

