



Uganda Russia Boston Alcohol Network for
Alcohol Research Collaboration on HIV/AIDS

URBAN ARCH Baseline Descriptive Data

ADEPTT – Uganda ARCH (n=167/300)	
<u>DEMOGRAPHICS</u>	N (%)
Age, mean (SD)	40 (8.0)
Male	84 (50.3)
Married	114 (68.3)
Basic education or higher*	42 (25.2)
Unemployed	6 (3.6)
<u>HEALTH INDICATORS</u>	
Depressive symptoms†	37 (22.2)
Antiretroviral medication, current	167 (100.0)
CD4 count, mean (SD)	715 (298)
HIV viral load suppressed, n=156	145 (92.9)
Hepatitis C infection (self-report)	N/A
Moderate/extreme pain interference, past 30 days	6 (3.6)
Experienced a fall, past 3 months	2 (1.2)
Broken bone from fall, past 3 months	1 (0.6)
TB, ever told by healthcare provider‡	1 (0.6)
Yes - Active	0 (0.0)
Yes - Latent	1 (0.6)
<u>SUBSTANCE USE</u>	
Alcohol use disorder, past year§	55 (33.1)
Hazardous alcohol use, past 3 months	65 (39.2)
IDU as HIV transmission route	0 (0.0)
Any illicit opioid use, lifetime	0 (0.0)
Any illicit opioid use, past 3 months	0 (0.0)
Marijuana use, past 3 months	0 (0.0)
Cocaine use, past 3 months	0 (0.0)
Current smoker	26 (15.6)

St. PETER – Russia ARCH (n= 267/400)	
<u>DEMOGRAPHICS</u>	N (%)
Age, mean (SD)	38.4 (6.3)
Male	179 (67.0)
Married or living with a partner	127 (47.6)
Basic education or higher*	258 (96.6)
Unemployed	94 (35.2)
<u>HEALTH INDICATORS</u>	
Depressive symptoms†	106 (39.8)
Antiretroviral medication, current	186 (69.7)
CD4 count, mean (SD)	411 (256)
HIV viral load suppressed (<300)	143 (57.2)
Hepatitis C infection (self-report)	207 (77.5)
Moderate/extreme pain interference, past 30 days	21 (7.9)
Experienced a fall, past 6 months	29 (10.9)
Broken bone from fall, past 6 months	1 (0.4)
TB, ever told by healthcare provider‡	41 (15.4)
Yes - Active	21 (7.9)
Yes - Latent	13 (4.9)
<u>SUBSTANCE USE</u>	
Alcohol use disorder, past year§	241 (90.2)
Heavy alcohol use, past 30 days ‡	265 (99.3)
IDU as HIV transmission route	202 (76.5)
Any illicit opioid use, lifetime	209 (78.3)
Any illicit opioid use, past 30 days	74 (27.7)
Marijuana use, past 30 days	34 (12.7)
Cocaine use, past 30 days	3 (1.1)
Current smoker‡	267 (100.0)

4F – Boston ARCH (n=163/400)	
<u>DEMOGRAPHICS</u>	N (%)
Age, mean (SD)	53.9 (9.4)
Male	99 (60.7)
Married or partnered	92 (56.8)
Basic education or higher*	114 (69.9)
Unemployed	132 (81.0)
<u>HEALTH INDICATORS</u>	
Depressive symptoms†	90 (55.2)
Antiretroviral medication, current	153 (93.9)
CD4 count, mean (SD)	676 (353)
HIV viral load suppressed (<200)	137 (88.4)
Hepatitis C infection (ever diagnosed)	91 (55.8)
Moderate/extreme pain interference, past 4 weeks	60 (36.8)
Experienced a fall, past 6 months	56 (34.4)
Broken bone from fall, past 6 months	3 (1.9)
TB, ever told by healthcare provider‡	19 (11.7)
Yes - Active	4 (2.5)
Yes - Latent	15 (9.2)
<u>SUBSTANCE USE</u>	
Alcohol use disorder, past year§	62 (38.0)
Heavy alcohol use, past 14 days	54 (33.1)
IDU as HIV transmission route	69 (42.6)
Any illicit opioid use, lifetime€	99 (67.8)
Any illicit opioid use, past 30 days	22 (13.5)
Marijuana use, past 30 days	71 (43.6)
Cocaine use, past 30 days	38 (23.3)
Current smoker	107 (65.6)

* ≥ 9 grades

† Based on CESD ≥ 16

‡ Via self-report. Latent TB is an eligibility criterion, so participants are confirmed positive with a TB skin test prior to enrolling

§ Based on DSM 5 criteria

|| Based on AUDIT-C

* ≥ 9 grades

† Based on CESD ≥ 16

‡ Via self-report. Study testing was not done.

§ Based on DSM 5 criteria

|| Based on TLFB, NIAAA consumption criteria

‡ Eligibility criteria at screening

* ≥ High school or GED

† Based on CESD ≥ 10

‡ Via self-report. Study testing was not done.

§ Based on DSM 5 criteria

|| Based on TLFB, NIAAA consumption criteria

€ ARCH Bone data