

# Supplementary Appendix

Elchanan Ben-Porath\*      Eddie Dekel†      Barton L. Lipman‡

September 2026

## 1 Proof of Lemma 2

For notational brevity, let

$$\pi(d) = \mathbb{E}_v \left[ \sum_i P_i(v \mid d) v_i \right]$$

and

$$U_i(d) = \mathbb{E}_v P_i(v \mid d) - c_i e_i(d).$$

Because  $d$  is a probability measure over dynamic mechanisms,  $\pi(d)$  and the  $U_i(d)$ 's are linear in  $d$  in the sense that

$$\pi(\alpha d + (1 - \alpha)d') = \alpha\pi(d) + (1 - \alpha)\pi(d'), \quad \forall \alpha \in [0, 1], \quad d, d' \in D.$$

Written this way,  $\pi(d)$  and the  $U_i(d)$ 's are convex and concave functions of  $d$ . This is because convex combinations of mixed strategies induce convex combinations of the distributions over outcomes.

So our problem is

$$\max_{d \in D} \pi(d)$$

subject to

$$U_i(d) \geq 0, \quad \forall i.$$

---

\*Department of Economics and Center for Rationality, Hebrew University. Email: benporat@math.huji.ac.il.

†Economics Department, Northwestern University, and School of Economics, Tel Aviv University. Email: eddiekel@gmail.com.

‡Department of Economics, Boston University. Email: blipman@bu.edu.

Let  $D^*$  denote the set of  $d$ 's solving this constrained optimization problem. It is not hard to use standard arguments (e.g., Theorem 2.1 in Balder (1989)) to show that the space of inducible distributions over outcomes is compact in the weak topology which, combined with the continuity of the utility functions, implies that  $D^* \neq \emptyset$ .

Let  $D^{**}(\lambda)$  denote the set of  $d$ 's solving

$$\max_{d \in D} \pi(d) + \sum_i \lambda_i U_i(d)$$

and let  $D^{**}$  denote the set of  $d$ 's such that there exists  $\lambda^* \in \mathbf{R}_+^N$  with  $d \in D^{**}(\lambda^*)$  such that (a)  $U_i(d) \geq 0$  for all  $i$  and (b)  $\lambda_i^* U_i(d) = 0$  for all  $i$ .

We now show that  $D^* = D^{**}$ .

First, we show  $D^{**} \subseteq D^*$ . Fix any  $d^* \in D^{**}$  and let  $\lambda^*$  be the associated vector in  $\mathbf{R}_+^N$ . Suppose, contrary to our claim, that there exists  $\hat{d}$  with  $U_i(\hat{d}) \geq 0$  for all  $i$  and  $\pi(\hat{d}) > \pi(d^*)$ . By  $d^* \in D^{**}(\lambda^*)$ ,

$$\pi(d^*) + \sum_{i=1}^N \lambda_i^* U_i(d^*) \geq \pi(\hat{d}) + \sum_{i=1}^N \lambda_i^* U_i(\hat{d}).$$

Because  $\lambda_i^* U_i(d^*) = 0$  for all  $i$ , this implies

$$\pi(d^*) \geq \pi(\hat{d}) + \sum_{i=1}^N \lambda_i^* U_i(\hat{d}).$$

Because  $\lambda_i^* \geq 0$  for all  $i$  and  $U_i(\hat{d}) \geq 0$  for all  $i$ , this implies  $\pi(d^*) \geq \pi(\hat{d})$ , a contradiction. Hence  $D^{**} \subseteq D^*$ .

The proof of the converse is a simplification of the proof of Theorem 1, Section 8.3, of Luenberger (1969). Fix any  $d^*$  in  $D^*$ . Let

$$A = \{u = (u_0, u_1, \dots, u_N) \in \mathbf{R}^{N+1} \mid \exists d \in D \text{ with } u_0 \leq \pi(d) \text{ and } u_i \leq U_i(d), \forall i = 1, \dots, N\}$$

$$B = \{u = (u_0, u_1, \dots, u_N) \in \mathbf{R}^{N+1} \mid u_0 \geq \pi(d^*) \text{ and } u_i \geq 0, \forall i = 1, \dots, N\}.$$

Obviously, both sets are nonempty as  $(\pi(d^*), 0, 0, \dots, 0)$  is in both sets.

Also, both sets are convex. The proof for  $B$  is trivial. For  $A$ , suppose  $u$  and  $u'$  are elements of  $A$  and fix any  $\alpha \in (0, 1)$ . Since  $u \in A$ , there exists  $d \in D$  satisfying

$$u_0 \leq \pi(d)$$

$$u_i \leq U_i(d), \forall i$$

and let  $d' \in D$  satisfy the analog for  $u'$ . Then we have

$$\alpha u_0 + (1 - \alpha)u'_0 \leq \alpha \pi(d) + (1 - \alpha)\pi(d') = \pi(\alpha d + (1 - \alpha)d')$$

and

$$\alpha u_i + (1 - \alpha)u'_i \leq \alpha U_i(d) + (1 - \alpha)U_i(d') = U_i(\alpha d + (1 - \alpha)d'),$$

implying  $\alpha u + (1 - \alpha)u' \in A$ .

Also, we have  $A \cap \text{int}(B) = \emptyset$ . To see this, suppose to the contrary that there is  $u \in \text{int}(B)$  with  $u \in A$ . Because  $u \in \text{int}(B)$ , we have  $u_0 > \pi(d^*)$  and  $u_i > 0$  for all  $i$ . Because  $u \in A$ , there exists  $d \in D$  with  $\pi(d) \geq u_0 > \pi(d^*)$  and  $U_i(d) \geq u_i > 0$  for all  $i$ . But this contradicts  $d^* \in D^*$  as  $d$  satisfies the constraints and gives a higher payoff than  $d^*$ .

By the Separating Hyperplane Theorem, there exists  $p \in \mathbf{R}^{N+1}$ ,  $p \neq 0$ , such that

$$p_0 u_0 + \sum_{i=1}^N p_i u_i \leq p_0 \hat{u}_0 + \sum_{i=1}^N p_i \hat{u}_i, \quad \forall u \in A, \hat{u} \in B.$$

We now show that  $p_i \geq 0$  for all  $i$ . Suppose to the contrary that some  $p_i < 0$ . Given the definition of  $B$ , we could make the corresponding component of  $\hat{u}$  arbitrarily large and violate this inequality, a contradiction.

Also,  $p_0 > 0$ . To see this, suppose that  $p_0 = 0$ . We know that  $(\pi(d^*), 0, \dots, 0) \in B$ , so this implies

$$\sum_{i=1}^N p_i u_i \leq 0,$$

for all  $u \in A$ . But consider the  $d \in D$  where we randomize uniformly over which agent to ask first and always give her the good. For this procedure,  $U_i(d) = (1 - c_i)/N > 0$  for all  $i$ . Hence there exists  $u \in A$  with  $u_i > 0$  for  $i = 1, \dots, N$ . Hence the only way this inequality could hold is if  $p_i = 0$  for all  $i$ . But we know  $p \neq 0$ , a contradiction.

For  $i = 1, \dots, N$ , let  $\lambda_i = p_i/p_0$ . Then we have  $\lambda \in \mathbf{R}_+^N$  with

$$u_0 + \sum_{i=1}^N \lambda_i u_i \leq \hat{u}_0 + \sum_{i=1}^N \lambda_i \hat{u}_i, \quad \forall u \in A, \hat{u} \in B.$$

Again,  $(\pi(d^*), 0, \dots, 0) \in B$ , so this implies

$$\pi(d^*) \geq u_0 + \sum_{i=1}^N \lambda_i u_i, \quad \forall u \in A.$$

For every  $d \in D$ ,  $(\pi(d), U_1(d), \dots, U_N(d)) \in A$ , so this implies

$$\pi(d^*) \geq \pi(d) + \sum_{i=1}^N \lambda_i U_i(d), \quad \forall d \in D.$$

In particular,  $d^* \in D$ , so this implies

$$\pi(d^*) \geq \pi(d^*) + \sum_i \lambda_i U_i(d^*).$$

Because  $\lambda_i \geq 0$  for all  $i$  and  $U_i(d^*) \geq 0$  for all  $i$ , we have  $\lambda_i U_i(d^*) = 0$  for all  $i$ . Hence

$$\pi(d^*) = \max_{d \in D} \left[ \pi(d) + \sum_{i=1}^N \lambda_i U_i(d) \right].$$

Rephrasing, this shows that there exists  $\lambda \in \mathbf{R}_+^N$  with  $d^* \in D^{**}(\lambda)$  with  $U_i(d^*) \geq 0$  and  $\lambda_i U_i(d^*) = 0$  for all  $i$ . Hence  $d^* \in D^{**}$ , completing the proof. ■

## 2 Border

In this section, we state and prove a version of a result in Border (1991). Lemma 1 below is essentially Border's Lemma 5.1 and Theorem 1 is essentially his Lemma 6.1.

First, we introduce some notation and terminology. In this section only, we denote the set of types for agent  $i$  by  $T_i$  and assume  $T_i$  is finite and not a singleton for each  $i$ . We consider *allocations*  $P = (P_1, \dots, P_N)$  with  $P_i : T \rightarrow [0, 1]$  with  $\sum_i P_i(t) \leq 1$  for all  $t \in T$ . Given  $P$ , we let  $p = (p_1, \dots, p_N)$  denote the *interim probabilities* where

$$p_i(t_i) = \sum_{t_{-i} \in T_{-i}} \mu_{-i}(t_{-i}) P_i(t_i, t_{-i}),$$

where  $\mu_j(t_j)$  is the (full support) prior over  $T_j$  and we assume type distributions are independent across agents. When  $p$  and  $P$  are related in this fashion, we say  $P$  *generates*  $p$ .

**Lemma 1.** *Any interim allocation  $p$  satisfies the following for every  $(\hat{T}_1, \dots, \hat{T}_N)$  with  $\hat{T}_i \subseteq T_i$  for all  $i$ :*

$$\sum_i \sum_{t_i \in \hat{T}_i} p_i(t_i) \mu_i(t_i) \leq 1 - \prod_i [1 - \mu_i(\hat{T}_i)].$$

*Proof.* The left-hand side is the probability that the good is allocated to some type in  $\cup_i \hat{T}_i$ . The right-hand side is the probability that at least one agent's type is in her  $\hat{T}_i$  set. ■

A *hierarchical allocation* is an allocation  $P$  that can be constructed as follows. We have a ranking function  $R$  which maps  $\cup_i T_i$  to  $\{1, \dots, K\}$  for some positive integer  $K$ . We assume that for every  $k < K$ , there is exactly one  $i$  such that  $R(t_i) = k$  for some  $t_i \in T_i$ . Note that this  $i$  may have more than one type with rank  $k$ . Note also that this restriction does *not* apply to rank  $K$  — there may be no or many agents with types at rank  $K$ .

Then given a type profile  $t = (t_1, \dots, t_N)$ , either all agents have rank  $K$  or there is a unique  $i$  with  $R(t_i) < R(t_j)$  for all  $j \neq i$ . If all agents have rank  $K$ , then  $P_j(t) = 0$  for all  $j$ . If there is a unique  $i$  with  $R(t_i) < R(t_j)$  for all  $j \neq i$ , then  $P_i(t) = 1$ . In other words, unless all agents are in the lowest rank, the agent who has the highest ranked type receives the good (where higher ranks have lower numbers).

We say that  $p$  is a *hierarchical interim probability* if it is generated by a hierarchical allocation  $P$ . Of course, the collection of hierarchical interim probabilities is a subset of the interim probabilities.

**Theorem 1.** *The set of hierarchical interim probabilities is the set of extreme points of the set of interim probabilities. That is, a function  $p$  is an interim probability if and only if it is a convex combination of hierarchical interim probabilities.*

*Proof.* We first show that any hierarchical interim probability  $p$  is an extreme point of the set of interim probabilities.

Fix a hierarchical interim allocation  $p$  and the ranking function  $R$  corresponding to the  $P$  that generates it. Given any rank  $k < K$ , let  $i(k)$  denote the unique agent  $i$  with a type  $t_i$  satisfying  $R(t_i) = k$  and let  $\hat{T}(k)$  denote the set of  $t_i \in T_{i(k)}$  with  $R(t_i) = k$ .

Suppose, contrary to what we wish to show, that  $p$  is not an extreme point of the set of interim probabilities. Then there exist interim probabilities  $q^1$  and  $q^2$ , neither equal to  $p$ , and  $\lambda \in (0, 1)$  such that  $\lambda q^1 + (1 - \lambda)q^2 = p$ . We obtain a contradiction by showing that we must have  $q^1 = q^2 = p$ .

Clearly, if  $K = 1$ , there is only one rank and all types of all agents have rank  $K$ . In this case,  $p$  is the zero vector, so the only interim probabilities  $q^1$  and  $q^2$  that could satisfy  $\lambda q^1 + (1 - \lambda)q^2 = p$  for  $\lambda \in (0, 1)$  are also the zero vector, establishing our claim.

So assume  $K \geq 2$ . Fix any  $t_{i(1)} \in \hat{T}(1)$ . Then  $p_{i(1)}(t_{i(1)}) = 1$ , so  $\lambda q^1 + (1 - \lambda)q^2 = p$  implies  $q^j_{i(1)}(t_{i(1)}) = 1$  for  $j = 1, 2$ .

This initiates an induction. Let  $K$  be the number of ranks. Suppose we have shown that for all  $k \leq \bar{k} < K$ , we have

$$q_{i(k)}^1(t_{i(k)}) = q_{i(k)}^2(t_{i(k)}) = p_{i(k)}(t_{i(k)}), \quad \forall t_{i(k)} \in \hat{T}(k).$$

We now show the same is true for rank  $k = \bar{k} + 1$ . This is obvious if  $\bar{k} + 1 = K$  since  $p_i(t_i) = 0$  for any  $t_i$  with rank  $K$ . So suppose  $\bar{k} + 1 < K$ . Let  $i = i(\bar{k} + 1)$  and fix any  $t_i^* \in \hat{T}(\bar{k} + 1)$ .

We have

$$p_{i(k)}(t_{i(k)}) = \Pr \left( \left\{ t_{-i(k)} \mid t_\ell \notin \hat{T}(j), \quad j = 1, \dots, k-1, \quad \forall \ell \neq i(k) \right\} \right), \quad \forall t_{i(k)} \in \hat{T}(k), \forall k \leq \bar{k}$$

and

$$p_i(t_i^*) = \Pr \left( \left\{ t_{-i} \mid t_{-i(k)} \notin \hat{T}(j), \quad j = 1, \dots, \bar{k}, \quad \forall \ell \neq i(k) \right\} \right).$$

Consider the inequality stated in Lemma 1 for the sets  $\hat{T}(k)$ ,  $k = 1, \dots, \bar{k}$ , and  $\{t_i^*\}$ . (If some agent  $j$  has no type in one of these sets, then  $\hat{T}_j = \emptyset$ . If some agent  $j$  has types in more than one of these sets, we use the union of the sets for  $j\hat{T}_j$ .) The left-hand side is

$$\sum_{k=1}^{\bar{k}} \sum_{t_{i(k)} \in \hat{T}(k)} p_{i(k)}(t_{i(k)}) \mu_{i(k)}(t_{i(k)}) + p_i(t_i^*) \mu_i(t_i^*).$$

Given the expression above for  $p_{i(k)}(t_{i(k)})$ , it is not hard to see that the first term is the probability that the highest-ranked agent has a rank of  $\bar{k}$  or higher. Equivalently, this is the probability that at least one agent has a rank of  $\bar{k}$  or higher. So the total probability is the probability that either one of the agents has a rank of  $\bar{k}$  or higher or else  $i$  is type  $t_i^*$ .

The right-hand side of the inequality is 1 minus the probability that no type is in one of these sets. That is, the right-hand side is

$$\leq 1 - \Pr(t_{i(k)} \notin \hat{T}(k), \quad k \leq \bar{k}, \quad \text{and} \quad t_i \neq t_i^*).$$

This must hold with equality. The first expression is exactly the probability that one of these types materializes, while the second is 1 minus the probability that none of them do.

Because the inequality holds with equality, we see that given the way we specified  $q^j$  on the types ranked above  $\bar{k}$ , we cannot set  $q_i^j(t_i^*) > p_i(t_i^*)$  for either  $j$  since doing so would give an interim probability that violates Lemma 1. Hence we again have  $q_i^j(t_i^*) = p_i(t_i^*)$  for  $j = 1, 2$ , completing the induction.

Hence every hierarchical interim probability is an extreme point of the set of hierarchical probabilities. Next, we show the converse: every extreme point of the set of interim probabilities is a hierarchical interim probability.

To show this, suppose not. Then there must be some interim probability, say  $p$ , which is not in the convex hull of the set of hierarchical interim probabilities. Let  $W$  denote this convex hull. Since  $W$  is convex and compact, there is a separating hyperplane  $f^*$ . In other words, viewing  $p$  and the elements of  $W$  as vectors, there exists a vector  $f^*$  such that  $f^* \cdot p > f^* \cdot q$  for all  $q \in W$ . Define  $f$  to be the vector with  $n$ th element  $f_n^*/\mu(n)$  where  $f_n^*$  is the  $n$ th element of  $f^*$  and  $\mu(n)$  is the probability of the type in the  $n$ th position in these vectors.

Without loss of generality, we can assume that the  $f_n$ 's are all distinct. That is, we have  $f_n \neq f_m$  for  $n \neq m$ . (If not, we can perturb  $f^*$  slightly to achieve this property.) Recall that the allocation that never gives the good to any agent is hierarchical. Hence the zero vector is contained in  $W$ . Hence  $f^* \cdot p > 0$  so  $f_n^* > 0$  for some  $n$  and hence  $f_n > 0$  for some  $n$ .

Without loss of generality, order the components of vectors so that  $f_1 > f_2 > \dots > f_M$  where  $M = \sum_i \#T_i$ . Hence we know that  $f_1 > 0$ . Hence there is some  $n^*$  with  $f_n > 0$  for  $n \leq n^*$  and  $f_n \leq 0$  for  $n \geq n^* + 1$  where  $n^*$  is the length of  $f$  if all components are positive.

We construct a hierarchical allocation and the associated  $q \in W$  as follows. Define the ranking  $R$  as follows. For  $n \leq n^*$ , assign rank  $n$  to the type in the  $n$ th component of these vectors. For every  $n \geq n^* + 1$ , assign rank  $K$  to the type in the  $n$ th component. Define functions  $i(k)$  and  $\hat{T}(k)$  for this ranking as above.

The corresponding  $q$  has 1 in the first component. The second component is

$$\Pr\left(\{t_{-i(2)} \mid t_j \notin \hat{T}(1), \forall j \neq i(2)\}\right),$$

the third is

$$\Pr\left(\{t_{-i(3)} \mid t_j \notin \hat{T}(1) \cup \hat{T}(2), \forall j \neq i(3)\}\right),$$

etc., and has 0 in all components from  $n^* + 1$  onward. We now show a contradiction to  $f^* \cdot p > f^* \cdot q$ .

We can write  $f^* \cdot p > f^* \cdot q$  as

$$\sum_{n=1}^M f_n \mu(n) p(n) > \sum_{n=1}^M f_n \mu(n) q(n) = \sum_{n=1}^{n^*} f_n \mu(n) q(n)$$

where  $p(n)$  is the  $n$ th component of the vector  $p$  and other terms are defined analogously.

Equivalently,

$$\sum_{n=1}^M f_n \mu(n)(p(n) - q(n)) > 0.$$

Since  $f_1 > 0$ , this implies

$$\sum_{n=2}^M \frac{f_n}{f_1} \mu(n)(p(n) - q(n)) > \mu(1)(q(1) - p(1)).$$

But  $q(1) = 1 \geq p(1)$ , so this implies

$$\sum_{n=2}^M \frac{f_n}{f_1} \mu(n)(p(n) - q(n)) > 0.$$

If  $f_2 \leq 0$ , this is a contradiction, since we would then have  $p(n) \geq 0 = q(n)$  and  $f_n \leq 0$  for all  $n \geq 2$ . So assume  $f_2 > 0$ .

By assumption,  $f_1/f_2 > 1$ . Hence

$$\frac{f_1}{f_2} \sum_{n=2}^M \frac{f_n}{f_1} \mu(n)(p(n) - q(n)) > \sum_{n=2}^M \frac{f_n}{f_1} \mu(n)(p(n) - q(n)) > \mu(1)(q(1) - p(1)).$$

That is,

$$\sum_{n=2}^M \frac{f_n}{f_2} \mu(n)(p(n) - q(n)) > \mu(1)(q(1) - p(1)),$$

so

$$\sum_{n=3}^M \frac{f_n}{f_2} \mu(n)(p(n) - q(n)) > \mu(2)(q(2) - p(2)) + \mu(1)(q(1) - p(1)).$$

It is not hard to see that the right-hand side must be non-negative. This follows from the fact that the inequality in Lemma 1 implies that  $\mu(1)q(1) + \mu(2)q(2)$  equals the maximum possible value for this sum. Hence  $\mu(1)p(1) + \mu(2)p(2)$  must be weakly smaller. Hence

$$\sum_{n=3}^M \frac{f_n}{f_2} \mu(n)(p(n) - q(n)) > 0.$$

Clearly, iterating, we obtain a contradiction. ■

**Remark 1.** Theorem 1 is slightly stronger than what we use. We only need the fact that every extreme point of the set of interim probabilities is a hierarchical interim probability, not the converse. We include the converse for the sake of completeness.

### 3 Completion of Proof of Lemma 6

In this section, we show for case (1) the alternative mechanism  $\bar{d}$  we constructed is incentive compatible and yields the principal a weakly higher expected payoff.

Because the distribution of  $\varphi(v_i)$  is the same as the distribution of  $v_j$ , the probability  $k$  is asked for evidence and/or receives the good is the same in  $d$  as in  $\bar{d}$  for all  $k \neq i, j$ . Also, the probability  $i$  gets the good in  $\bar{d}$  is the same as the probability  $j$  gets the good in  $d$  and the probability  $i$  is asked for evidence in  $\bar{d}$  is the same as the probability  $j$  is asked in  $d$ . Because  $c_i \leq c_j$ , then,  $i$ 's incentive constraint is satisfied. (The incentive constraint for  $j$  is trivially satisfied in  $\bar{d}$  as she is never asked for evidence.) Hence  $\bar{d}$  is incentive compatible.

So consider the principal's payoff from  $\bar{d}$ . Because the principal never gives the good to  $j$  in  $\bar{d}$  and because the probability the principal gives the good to any agent  $k \neq i, j$  is unchanged, we can write the payoff as

$$E_v[P_i(v_i, v_j, v_{-ij} \mid \bar{d})v_i] + \sum_{k \neq i, j} E_v[P_k(v \mid d)v_k].$$

By construction,  $i$ 's probability of receiving the good in  $\bar{d}$  when her type is  $v_i$  and the types of the agents other than  $i$  and  $j$  are  $v_{-ij}$  is independent of  $v_j$  (as  $j$  never provides evidence) and is the same as  $j$ 's probability of getting the good in  $d$  when  $j$ 's type is  $\varphi(v_i)$  and the types of the other agents are  $v_{-ij}$ , again for any type  $i$  might be. That is,

$$P_i(v_i, v'_j, v_{-ij} \mid \bar{d}) = P_j(v'_i, \varphi(v_i), v_{-ij} \mid d), \quad \forall v_i, v_{-ij}, v'_j, v'_i.$$

Also, the fact that the distribution of  $v_j$  is the same as the distribution of  $\varphi(v_i)$  implies that

$$E_v[P_j(v'_i, v_j, v_{-ij} \mid d)v_j] = E_v[P_j(v'_i, \varphi(v_i), v_{-ij} \mid d)\varphi(v_i)], \quad \forall v'_i.$$

The principal's expected payoff in  $\bar{d}$  is weakly larger than the expected payoff in  $d$  iff

$$E_v[P_i(v_i, v'_j, v_{-ij} \mid \bar{d})v_i] \geq E_v[P_j(v'_i, v_j, v_{-ij} \mid d)v_j]$$

or, using the above,

$$E_v[P_j(v'_i, \varphi(v_i), v_{-ij} \mid d)v_i] \geq E_v[P_j(v'_i, \varphi(v_i), v_{-ij} \mid d)\varphi(v_i)].$$

Hence the new mechanism yields the principal a weakly better payoff if  $v_i \geq \varphi(v_i)$  for all  $v_i$ , or  $v_i \geq F_j^{-1}(F_i(v_i))$  or  $F_j(v) \geq F_i(v)$  for all  $v$ . This holds as  $F_i$  FOSD  $F_j$ .

## References

- [1] Balder, E., “On Compactness of the Space of Policies in Stochastic Dynamic Programming,” *Stochastic Processes and their Applications*, **32**, June 1989, 141–150.