

Review

Psychotherapy for Military-Related PTSD

A Review of Randomized Clinical Trials

Maria M. Steenkamp, PhD; Brett T. Litz, PhD; Charles W. Hoge, MD; Charles R. Marmar, MD

IMPORTANCE Posttraumatic stress disorder (PTSD) is a disabling psychiatric disorder common among military personnel and veterans. First-line psychotherapies most often recommended for PTSD consist mainly of “trauma-focused” psychotherapies that involve focusing on details of the trauma or associated cognitive and emotional effects.

OBJECTIVE To examine the effectiveness of psychotherapies for PTSD in military and veteran populations.

EVIDENCE REVIEW PubMed, PsycINFO, and PILOTS were searched for randomized clinical trials (RCTs) of individual and group psychotherapies for PTSD in military personnel and veterans, published from January 1980 to March 1, 2015. We also searched reference lists of articles, selected reviews, and meta-analyses. Of 891 publications initially identified, 36 were included.

FINDINGS Two trauma-focused therapies, cognitive processing therapy (CPT) and prolonged exposure, have been the most frequently studied psychotherapies for military-related PTSD. Five RCTs of CPT (that included 481 patients) and 4 RCTs of prolonged exposure (that included 402 patients) met inclusion criteria. Focusing on intent-to-treat outcomes, within-group posttreatment effect sizes for CPT and prolonged exposure were large (Cohen *d* range, 0.78-1.10). CPT and prolonged exposure also outperformed waitlist and treatment-as-usual control conditions. Forty-nine percent to 70% of participants receiving CPT and prolonged exposure attained clinically meaningful symptom improvement (defined as a 10- to 12-point decrease in interviewer-assessed or self-reported symptoms). However, mean posttreatment scores for CPT and prolonged exposure remained at or above clinical criteria for PTSD, and approximately two-thirds of patients receiving CPT or prolonged exposure retained their PTSD diagnosis after treatment (range, 60%-72%). CPT and prolonged exposure were marginally superior compared with non-trauma-focused psychotherapy comparison conditions.

CONCLUSIONS AND RELEVANCE In military and veteran populations, trials of the first-line trauma-focused interventions CPT and prolonged exposure have shown clinically meaningful improvements for many patients with PTSD. However, nonresponse rates have been high, many patients continue to have symptoms, and trauma-focused interventions show marginally superior results compared with active control conditions. There is a need for improvement in existing PTSD treatments and for development and testing of novel evidence-based treatments, both trauma-focused and non-trauma-focused.

JAMA. 2015;314(5):489-500. doi:10.1001/jama.2015.8370

← Editorial page 453

+ Author Audio Interview at jama.com

← Related article page 501

+ Supplemental content at jama.com

Author Affiliations: Steven and Alexandra Cohen Veterans Center for Posttraumatic Stress and Traumatic Brain Injury, New York University Langone School of Medicine, New York, New York (Steenkamp, Marmar); VA Boston Healthcare System, Massachusetts Veterans Epidemiological Research and Information Center (MAVERIC), Boston University School of Medicine, Boston (Litz); Walter Reed Army Institute of Research, Silver Spring, Maryland (Hoge).

Corresponding Author: Maria M. Steenkamp, PhD, Steven and Alexandra Cohen Veterans Center for Posttraumatic Stress and Traumatic Brain Injury, New York University Langone School of Medicine, One Park Ave, Room 8-107, New York, NY 10016 (maria.steenkamp@nyumc.org).

Section Editor: Mary McGrae McDermott, MD, Senior Editor.

Posttraumatic stress disorder (PTSD) is a disabling psychiatric condition common among military personnel and veterans, and consequently, a significant public health challenge. Approximately 13% of Iraq or Afghanistan veterans¹ and 10% of Gulf War veterans who experienced combat have PTSD,² and 11% of Vietnam veterans continue to report PTSD symptoms that impair functioning 40 years after the war.³ Military-related PTSD is often accompanied by a variety of mental and physical health conditions, particularly depression, anxiety, and substance misuse.^{4,5} War veterans with PTSD also have extensive functional impairments such as unemployment and income disparities,⁶ family and relationship difficulties,⁷ aggressive behavior,⁸ and poor quality of life.⁹ Twenty-three percent of Vietnam veterans with PTSD (compared with 4% among those without PTSD) reported being unemployed when assessed 15 or more years after service, 33% (compared with 16%) reported perpetrating serious interpersonal violence in the past year, and 40% (compared with 10%) reported low well-being.¹⁰ Risk factors for PTSD in military populations include war zone exposure, being wounded, younger age when deployed, less education, greater exposure to childhood trauma, and less social support during and after deployment.¹¹ If left untreated, military-related PTSD often follows a chronic course, resulting in lifelong dysfunction.¹²

Over the past 10 years, an increasing number of randomized clinical trials (RCTs) of PTSD treatments in military personnel and veterans have emerged, coinciding with a major policy shift in the Departments of Defense (DoD) and Veterans Affairs (VA) toward therapies considered evidence-based.¹³ Psychotherapy is more consistently recommended as first-line treatment for PTSD than medications across clinical practice guidelines and in DoD and VA practice settings. In this review, we focus on RCTs of individual and group psychotherapies for PTSD in military and veteran populations to examine the degree of symptom improvement and efficacy relative to control conditions.

Methods

We searched PubMed, PsycINFO, and PILOTS for RCTs of psychotherapy for PTSD among military personnel or veterans, published from January 1980 (the year the PTSD diagnosis was first introduced) to March 1, 2015. PTSD was defined according to the diagnostic criteria accepted at the time of the RCTs, which most often followed the *Diagnostic and Statistical Manual for Mental Disorders* (Fourth Edition, Text Revision).¹⁴ This definition included 3 clusters of symptoms following trauma, present for at least 1 month, including reexperiencing the trauma (eg, intrusive thoughts or nightmares), avoiding reminders of the trauma, and hyperarousal (eg, hypervigilance, irritability, difficulty concentrating). We also manually searched reference lists of articles, selected reviews, and meta-analyses.

We selected only English-language RCTs that (1) were conducted with service members, veterans, or both; (2) reported PTSD as an inclusion criterion; (3) reported at least pretreatment and post-treatment total scores on standardized PTSD clinical measures; (4) used either individual or group psychotherapy; and (5) did not solely involve pharmacotherapies, pharmacologically augmented psychotherapy, or other biological treatments. We did not include

trials of collaborative care models, dosing studies, trials targeting specific symptoms (eg, insomnia, anger) rather than the full syndrome, or trials targeting substance use disorders comorbid with PTSD. We focused on intent-to-treat outcomes, when available.

Results

Of 891 publications initially identified, 36 were included (Table 1^{15-27,29,58} and Table 2³⁰⁻⁵⁰; eFigure in the Supplement). We grouped trials of the most commonly studied first-line trauma-focused therapies (ie, therapies given the highest evidence recommendations in clinical guidelines) (Table 1), followed by second-line interventions (ie, therapies for which there is less evidence supporting effectiveness) (Table 2). The principal efficacy criteria, reported variably in published trials, included degree of clinically significant PTSD symptom improvement (typically defined as a 10- or 12-point decline in self-reported or interviewer-assessed PTSD symptoms),²³ mean PTSD symptom level at posttreatment and follow-up, loss of PTSD diagnosis, degree of symptom remission, and effect sizes (most commonly Cohen *d*, calculated as the difference between 2 mean PTSD severity scores divided by the pooled SD [a *d* of 0.20 indicates a small effect size; a *d* of 0.50, a medium effect size; and a *d* of 0.80, a large effect size]).

First-Line Psychotherapies

The diverse range of PTSD psychotherapies are broadly grouped into "trauma-focused" and non-trauma-focused. Trauma-focused therapies are cognitive-behavioral treatments that involve a range of techniques that attend to the details of the trauma or associated emotions or cognitive processes (beliefs, assumptions). The 3 most widely studied trauma-focused therapies, which are considered leading evidence-based psychotherapies by all major clinical guidelines,⁵¹ are cognitive processing therapy (CPT), prolonged exposure therapy, and eye movement desensitization and reprocessing (EMDR) therapy (Box). All are manualized (ie, protocolized in a session-by-session manner), delivered principally in specialty care settings, use different techniques and theoretical rationales, require sustained engagement (typically 12 sessions), and can be emotionally demanding for patients. Prolonged exposure includes asking patients to repeatedly recount the trauma to extinguish fear responses associated with the memory (a technique known as imaginal exposure) and to practice facing trauma reminders and triggers in the real world (known as in vivo exposure). CPT involves changing maladaptive beliefs related to the trauma (known as cognitive restructuring), with the option of writing an account of the trauma. EMDR also comprises exposure and cognitive restructuring elements but asks patients to maintain dual focus on an external stimulus (eg, eye-movement tracking of therapist hand movements) while thinking about the trauma.

Meta-analyses of mostly civilian studies show large pre-post treatment effects (within-group) and between-group effects compared with control conditions for these treatments, with generally comparable outcomes for CPT, prolonged exposure, EMDR, and other trauma-focused modalities.⁵² In contrast, non-trauma-focused therapies attend principally to current life stressors, reactions, goals, or relationships. The only non-trauma-focused therapy that has received high-level evidence statements in PTSD clinical

Table 1. Summary of Outcomes From RCTs of First-Line Interventions for Military-Related PTSD

Source	Participants (No.)	Conditions	Longest Follow-up, mo	No. (%)		Remission at Post-Treatment	Mean Posttreatment Total PTSD Score	Pre-Post Decrease in PTSD Symptoms	No. (%)	% Attaining Clinically Meaningful Change at Posttreatment	% Retaining PTSD Diagnosis at Posttreatment	Between-Group ES (95% CI) From Pretreatment to Posttreatment
				Drop Out	Drop Out							
CPT												
Monson et al, ¹⁵ 2006	Veterans (30)	CPT	1	6 (20)	NR	NR	52.14 (CAPS IIT)	24.59 (CAPS IIT)	15 (50) (CAPS IIT)	18 (60) (CAPS IIT)	18 (60) (CAPS IIT)	$g = 1.12$ (-0.58 to 1.67) (CAPS IIT)
	Veterans (30)	Waitlist	1	4 (13)	NR	NR	76.03 (CAPS IIT)	3.07 (CAPS IIT)	3 (10) (CAPS IIT)	29 (97) (CAPS IIT)	29 (97) (CAPS IIT)	
Forbes et al, ¹⁶ 2012	Australian veterans (30)	CPT	3	9 (30)	6 (27) (CAPS IIT)	NR	48.03 (CAPS IIT)	27.5 (CAPS IIT)	16 (67) (CAPS IIT)	15 (63) (CAPS completer)	15 (63) (CAPS completer)	$d = 0.97$ (0.43 to 1.51)
	Australian veterans (29)	TAU	3	9 (31)	1 (3) (CAPS IIT)	NR	57.73 (CAPS IIT)	6.82 (CAPS IIT)	8 (35) (CAPS IIT)	20 (87) (CAPS completer)	20 (87) (CAPS completer)	
Surís et al, ¹⁷ 2013	Veterans with military sexual trauma (72)	CPT	6	18 (35)	NR	NR	64.97 (CAPS IIT)	20.1 (CAPS IIT)	66% (CAPS IIT)	72% (CAPS IIT)	72% (CAPS IIT)	$d = 0.49$ (0.22 to -0.76) (CAPS IIT)
	Veterans with military sexual trauma (57)	PCT	6	6 (18)	NR	NR	68.64 (CAPS IIT)	15.17 (CAPS IIT)	48% (CAPS IIT)	74% (CAPS IIT)	74% (CAPS IIT)	
Morland et al, ¹⁸ 2014	Veterans (61)	Group CPT-C via telemedicine	6	10 (16)	NR	NR	58.7 (CAPS IIT)	10.2 (CAPS IIT)	71 (57) overall (CAPS IIT)	89 (71) overall (CAPS IIT)	89 (71) overall (CAPS IIT)	$d = 0.27$ (CAPS IIT)
	Veterans (64)	Group CPT-C in person	6	8 (13)	NR	NR	55.6 (CAPS IIT)	16.4 (CAPS IIT)	71 (57) overall (CAPS IIT)	89 (71) overall (CAPS IIT)	89 (71) overall (CAPS IIT)	
Resick et al, ¹⁹ 2015	Active duty soldiers (56)	Group CPT-C	12	15 (27)	NR	NR	23.0 (PSSI)	4.7 (PSSI)	15 (49) (PCL)	NR	NR	$d = 0.21$ (PSSI)
	Active duty soldiers (52)	Group PCT	12	7 (13)	NR	NR	23.9 (PSSI)	3.2 (PSSI)	14 (34) (PCL)	NR	NR	
Prolonged Exposure, Imaginal Exposure, and Group Exposure												
Cooper & Klum, ²⁰ 1989	Veterans (8)	TAU + implosive therapy/flooding	3	6 (27) overall	NR	NR	NR	NR	NR	NR	NR	NR
	Veterans (8)	TAU	3	6 (27) overall	NR	NR	NR	NR	NR	NR	NR	NR
Keane et al, ²¹ 1989	Vietnam veterans (11)	Implosive therapy	6	NR	NR	NR	NR	NR	NR	NR	NR	NR
	Vietnam veterans (13)	Waitlist	6	NR	NR	NR	NR	NR	NR	NR	NR	NR
Boudewyns & Hyer, ²² 1990	Inpatient veterans (19)	Exposure	3	13 (25) overall	NR	NR	NR	NR	NR	NR	NR	NR
	Inpatient veterans (19)	TAU	3	13 (25) overall	NR	NR	NR	NR	NR	NR	NR	NR
Schnurr et al, ²³ 2003	Veterans (180)	Group exposure	12	41 (23)	NR	NR	74.00 (CAPS IIT)	6.41 (CAPS IIT)	39% (CAPS IIT)	NR	NR	NR
	Veterans (180)	Group PCT	12	16 (9)	NR	NR	76.03 (CAPS IIT)	5.98 (CAPS IIT)	38% (CAPS IIT)	NR	NR	NR

(continued)

Table 1. Summary of Outcomes From RCTs of First-Line Interventions for Military-Related PTSD (continued)

Source	Participants (No.)	Conditions	Longest Follow-up, mo	Drop Out	Remission at Post-Treatment	Mean Posttreatment Total PTSD Score (CAPS ITT)	Pre-Post PTSD Symptoms	No. (%)	% Attaining Clinically Meaningful Change at Posttreatment (CAPS ITT)	% Retaining PTSD Diagnosis at Posttreatment (CAPS ITT)	Between-Group ES (95% CI) From Pretreatment to Posttreatment
Schnurr et al, ²⁴ 2007	Female veterans primarily with sexual trauma (141)	PE	6	53 (38)	24 (17)	52.9 (47.7 to 58.0) (CAPS ITT)	24.7 (CAPS ITT)	99 (70) (CAPS ITT)	86 (61) (CAPS ITT)		
	Female veterans primarily with sexual trauma (143)	PCT	6	30 (21)	10 (7)	60.1 (55.3 to 64.8) (CAPS ITT)	17.8 (CAPS ITT)	84 (59) (CAPS ITT)	114 (80) (CAPS ITT)		<i>d</i> = 0.27 CAPS ITT
Nacasch et al, ²⁵ 2011	Israeli survivors of combat or terrorism (15)	PE	12	2 (13)	NR	18.9 (PSSI ITT)	18.2 (PSSI ITT)	NR	NR		
	Israeli survivors of combat or terrorism (15)	TAU	12	2 (13)	NR	35.0 (PSSI ITT)	1.8 (PSSI ITT)	NR	NR		<i>d</i> = 1.80 PSSI ITT
Yehuda et al, ²⁶ 2014	Veterans (37)	PE	3	12 (31)	NR	NR	23 (CAPS)	NR	14 (56) (CAPS completer)		NR
	Veterans (15)	Minimal attention	3	3 (21)	NR	NR	14 (CAPS)	NR	10 (83) (CAPS completer)		NR
Rauch et al, ²⁷ 2015	Veterans (18)	PE	None	7 (39)	NR	30.0 (CAPS completer)	49.2 (CAPS completer)	10 (91) (CAPS completer)	NR		<i>d</i> = 0.98 CAPS completer
	Veterans (18)	PCT	None	3 (17)	NR	53.6 (CAPS completer)	23.8 (CAPS completer)	9 (60) (CAPS completer)	NR		
EMDR											
Boudewyns & Hyer, ⁵⁸ 1996	Veterans (21)	EMDR	None	NR	NR	50.09 (CAPS)	25.14 (CAPS)	NR	NR		
	Veterans (18)	EMDR without eye movement	None	NR	NR	67.50 (CAPS)	18.1 (CAPS)	NR	NR		NR
	Veterans (22)	TAU	None	NR	NR	67.18 (CAPS)	14.05 (CAPS)	NR	NR		NR
Carlson et al, ²⁹ 1998	Veterans (10)	EMDR	3	0	NR	NR	NR	NR	2 (22)		
	Veterans (13)	Biofeedback relaxation	3	1 (3)	NR	NR	NR	NR	7 (78)		
	Veterans (12)	Routine care	3	0	NR	NR	NR	NR	NR		NR
				12 (26) overall prior to group assignment							

Abbreviations: CAPS, Clinician Administered PTSD Scale; completer, treatment completer sample only; therapy, PSSI, PTSD Symptom Scale Interview; PTSD, posttraumatic stress disorder; RCT, randomized clinical trial; CPT, cognitive processing therapy; CPT-C, CPT cognitive-only version; EMDR, eye movement desensitization and reprocessing; ES, effect size; ITT, intent-to-treat; NR, not reported; PCL, PTSD Checklist; PCT, present-centered

practice guidelines is stress inoculation training, which involves stress management skills (eg, breathing, muscle relaxation) and applying these skills to day-to-day stressors and reminders of the trauma.

In 2008, CPT and prolonged exposure were selected by the VA for nationwide dissemination in an attempt to better standardize care for veterans, and 98% of VA centers now offer both.^{53,54} Initially, neither intervention was validated sufficiently in active duty military or veteran populations; they were originally tested largely in civilian female survivors of sexual assault. EMDR has not been disseminated within the VA and DoD, and EMDR research has received comparatively little VA or DoD funding but has strong evidence in civilian studies and high-level endorsement from many guidelines internationally. RCTs of stress inoculation training are lacking in veterans, and it is infrequently used in the VA and DoD.

Efficacy of CPT

Five trials of CPT (that included 481 patients) met inclusion criteria; 4 enrolled veterans¹⁵⁻¹⁸ and 1 enrolled active-duty soldiers.¹⁹ The first included a waitlist comparison,¹⁵ and subsequent studies used either a treatment-as-usual comparison or an active comparison condition known as present-centered therapy, a non-trauma-focused treatment protocol focused largely on current life problems. One noninferiority trial compared group CPT delivered in person vs via telemedicine.¹⁸ Treatment dropout rates ranged from 16% to 35%.

Within-group intent-to-treat effect sizes for CPT were reported in 3 trials and were large ($d = 0.78$,¹⁸ $d = 1.10$,¹⁹ and $d = 1.02$ ¹⁷). All trials reported the percentage of patients attaining meaningful symptom change (49%-67%), and the mean posttreatment PTSD scores, which remained at or above clinical cutoffs (eg, 50 on the Clinician Administered PTSD Scale [CAPS] and 20 on the PTSD Scale-Interview [PSSI]), with a range of 48.03 to 64.97 for CAPS and 23.00 for PSSI. Four trials reported posttreatment PTSD diagnosis,¹⁵⁻¹⁸ with approximately two-thirds of patients retaining their diagnosis (60%-72%). Only 1 trial reported symptom remission (CAPS score <20) of 27%.¹⁶

Concerning comparisons to control conditions, CPT outperformed waitlist approaches for mean PTSD symptom reduction ($g = 1.12$), loss of diagnosis, and meaningful symptom change, although the latter was no longer statistically significant by 1 month posttreatment.¹⁵ CPT produced significantly greater symptom remission and clinically meaningful symptom improvement than treatment as usual in a study of Australian veterans ($d = 0.97$). Group differences in loss of diagnosis and meaningful symptom improvement were nonsignificant at 3-month follow-up but remained significant for symptom remission.¹⁶ In a trial comparing CPT and present-centered therapy for military sexual trauma, CPT resulted in significantly greater reduction in self-reported PTSD symptoms at posttreatment ($d = 0.85$), although group differences were no longer significant by 2-, 4-, or 6-month follow-up.¹⁷ There were no significant group differences in interviewer-assessed PTSD symptoms. A trial comparing group CPT with group present-centered therapy in active-duty soldiers also found that self-reported PTSD symptoms improved in both groups but statistically significantly more so in the CPT group ($d = 0.40$).¹⁹ Between-group differences were small and not significant for interviewer-assessed PTSD symptoms at posttreatment ($d = 0.21$), 6-month ($d = 0.22$), and 12-month ($d = 0.21$) follow-up. There were no significant differences between groups in

the percentage of patients attaining clinically significant change in self-reported symptoms at posttreatment, 6-month, or 12-month follow-up.

In sum, trials of CPT for military-related PTSD have included both veterans and active-duty personnel with combat or military sexual trauma, have shown high methodological rigor (although fidelity problems were present in 1 trial),¹⁷ and have had large effect sizes when compared with no treatment (waitlist) or treatment as usual. However, CPT was marginally superior to active, non-trauma-focused control comparisons.

Efficacy of Prolonged Exposure

Four RCTs of prolonged exposure (that included 402 patients) met inclusion criteria.²⁴⁻²⁷ Earlier trials examined similar exposure-based mechanisms but not the full prolonged exposure protocol²⁰⁻²² and included a large trial of an exposure-based group therapy that did not lead to meaningful PTSD symptom reduction or outperform a present-centered control.²³ The most robust trial of prolonged exposure compared prolonged exposure with present-centered therapy in female veterans with sexual trauma,²⁴ while 3 small studies of combat veterans used minimal attention,²⁶ treatment as usual,²⁵ or present-centered therapy²⁷ control conditions. Two of these trials^{26,27} focused primarily on glucocorticoid-related biomarker responses associated with clinical improvement. Treatment dropout rates ranged from 13% to 39%.

Within-group intent-to-treat effect size for prolonged exposure was reported in 1 trial and was large ($d = 0.80$).²⁴ One trial reported on clinically meaningful PTSD symptom reduction in the intent-to-treat sample, which occurred in 70% of patients.²⁴ Mean posttreatment intent-to-treat symptom scores were reported in 2 trials and remained at or above clinical cutoffs for PTSD (52.9 [CAPS]²⁴ and 18.9 [PSSI]²⁵). One trial reported loss of diagnosis in the intent-to-treat sample and found that 61% retained their diagnosis after treatment.²⁴ The only trial to report remission (≤ 20 CAPS) found remission in 17% of the intent-to-treat sample.²⁴

When compared with control conditions, both prolonged exposure and present-centered therapy improved symptoms in female veterans with predominantly sexual assault trauma, with a small intent-to-treat effect size favoring prolonged exposure ($d = 0.27$); there were no significant group differences for clinically meaningful improvement at any point and no significant group differences for loss of diagnosis or symptom remission at either the 3- or 6-month follow-up.²⁴ In a small sample of Israeli veterans, prolonged exposure resulted in greater symptom reduction than psychodynamic therapy ($d = 1.80$), and outcomes were maintained through 12-month follow-up.²⁵ Prolonged exposure failed to outperform a minimal-attention control that consisted of 30-minute weekly symptom monitoring phone calls; both groups significantly improved, with no significant group differences.²⁶ Last, a small RCT examining cortisol response following prolonged exposure found no significant differences between prolonged exposure and present-centered therapy on interviewer-assessed PTSD outcomes in the intent-to-treat sample, although significant differences favoring prolonged exposure were found in completers ($d = 3.16$ for prolonged exposure vs $d = 1.08$ for present-centered therapy).²⁷

In sum, fewer methodologically robust trials for military-related PTSD are available for prolonged exposure than for CPT. With

Table 2. Summary of Outcomes From RCTs of Second-Line Interventions for Military-Related PTSD

Source	Participants (No.)	Conditions	Longest Follow-up, mo	Drop Out	Remission at Post-Treatment	Mean Post-Treatment Total PTSD Score	Pre-Post Drop in PTSD Symptoms	No. (%)		Between-Group ES From Pretreatment to Posttreatment
								% Attaining Clinically Meaningful Change At Posttreatment	% Retaining PTSD Diagnosis at Posttreatment	
Watson et al, ³⁰ 1997	Vietnam veterans (30)	Relaxation	None	NR	NR	95.0 (PTSD-I completer)	0.4 (PTSD-I completer)	NR	NR	NR
	Vietnam veterans (30)	Relaxation + deep breathing	None	NR	NR	97.8 (PTSD-I completer)	0.3 (PTSD-I completer)	NR	NR	NR
	Vietnam veterans (30)	Relaxation + deep breathing + biofeedback	None	NR	NR	89.4 (PTSD-I completer)	1.1 (PTSD-I completer)	NR	NR	NR
Dunn et al, ³¹ 2007	Veterans (55)	Self-management therapy	12	17 (33)	NR	73.93 (CAPS completer)	2.01 (CAPS completer)	NR	NR	$d = 0.23$
	Veterans (56)	Psychoeducation group therapy	12	6 (12)	NR	77.10 (CAPS completer)	+1.05 (CAPS completer)	NR	NR	
Frieh et al, ³² 2007	Veterans (21)	In-person group CBT	3	9 (43)	NR	56.58 (PCL completer)	5.8 (PCL completer)	NR	NR	NR
	Veterans (17)	Telepsychiatry group CBT	3	8 (47)	NR	68.11 (PCL completer)	+1.11 (PCL completer)	NR	NR	NR
Litz et al, ³³ 2007	Active duty personnel (24)	Internet CBT	6	12 (27) overall	7 (42.9) (PSS-I <6)	14.86 (PSS-I completer)	9.16 (PSS completer)	NR	NR	$d = 0.41$
	Active duty personnel (21)	Internet supportive counseling	6	12 (27) overall	1 (6.3) (PSS-I <6)	20.00 (PSS-I completer)	11.85 (PSS-I completer)	NR	NR	
Bormann et al, ³⁴ 2008	Veterans (14)	Mantram repetition	None	4 (12) overall	NR	NR	4.79 (CAPS ITT)	NR	NR	$d = 0.33$
	Veterans (15)	Waitlist	None	NR	NR	NR	2.64 (CAPS ITT)	NR	NR	
Lande et al, ³⁵ 2010	Active duty soldiers (22)	Biofeedback + TAU	None	NR	NR	NR	NR	NR	NR	NR
	Active duty soldiers (17)	TAU	None	NR	NR	NR	NR	NR	NR	NR
Ready et al, ³⁶ 2010	Vietnam veterans (6)	Virtual reality exposure	6	1 (17)	NR	59.20 (CAPS completer)	31.8 (CAPS completer)	NR	NR	$d = 0.28$
	Vietnam veterans (5)	PCT	6	1 (20)	NR	75.50 (CAPS completer)	23.0 (CAPS completer)	NR	NR	
Beidel et al, ³⁷ 2011	Vietnam veterans (14)	Trauma management therapy	None	5 (14) overall	NR	69.0 (CAPS completer)	15.9 (CAPS completer)	NR	NR	NR
	Vietnam veterans (16)	Exposure therapy	None	NR	NR	65.5 (CAPS completer)	25.1 (CAPS completer)	NR	NR	NR
Kent et al, ³⁸ 2011	Veterans (20)	Resilience-oriented group therapy	None	1 (5)	NR	23.00 (PDS ITT)	12.90 (PDS ITT)	NR	NR	$d = 1.40$
	Veterans (19)	Waitlist	None	2 (11)	NR	36.90 (PDS ITT)	0.63 (PDS ITT)	NR	NR	

(continued)

Table 2. Summary of Outcomes From RCTs of Second-Line Interventions for Military-Related PTSD (continued)

Source	Participants (No.)	Conditions	Longest Follow-up, mo	Drop Out	Remission at Post-Treatment	Mean Post-Treatment Total PTSD Score	Pre-Post Drop in PTSD Symptoms	No. (%)		Between-Group ES From Pretreatment to Posttreatment
								% Attaining Clinically Meaningful Change At Posttreatment	% Retaining PTSD Diagnosis at Posttreatment	
McLay et al, ³⁹ 2011	Active duty service members (10)	Virtual reality graded exposure therapy	None	0	NR	48.1 (CAPS ITT)	35.4 (CAPS)	7 (70) showed improvement of >30%	NR	NR
Possemato et al, ⁴⁰ 2011	Active duty service members (10)	TAU	None	1 (10)	NR	72.3 (CAPS ITT)	9.4 (CAPS)	1 (11%)	NR	NR
Jain et al, ⁴¹ 2013	Active duty Marines (68)	Written emotional disclosure	3	0	NR	NR	NR	NR	NR	NR
Jain et al, ⁴¹ 2013	Active duty Marines (68)	HT + GI + TAU	None	6 (9)	NR	40.7 (37.0-44.2) (PCL ITT)	13.3 (PCL ITT)	NR	NR	NR
Kearney et al, ⁴² 2012	Veterans (22)	MBSR + TAU	4	2 (8)	NR	52.45 (PCL ITT)	7.43 (PCL ITT)	8 (36)	NR	d = 0.51 (0.11-1.12)
Niles et al, ⁴³ 2012	Veterans (17)	Mindfulness	6 wk	4 (24)	NR	47.46 (CAPS completer)	13.46 (CAPS completer)	5 (38.5) (20 points on CAPS)	NR	NR
Strachan et al, ⁴⁴ 2012	OEF/OIF veterans and active duty service members (20)	BA-TE in person	None	5 (25)	NR	47.9 (PCL completer)	11.1 (PCL completer)	NR	NR	NR
Bormann et al, ⁴⁵ 2013	Veterans (71)	Mantram repetition + TAU	6 wk	5 (7)	$\eta^2_p = .03$	66.16 (CAPS ITT)	16.92 (CAPS ITT)	18 (24)	NR	NR
Church et al, ⁴⁶ 2013	Veterans (30)	EFT + TAU	6	1 (3)	39 (80) overall (including the waitlist that received EFT)	72.59 (CAPS ITT)	10.24 (CAPS ITT)	9 (12)	NR	NR
	Veterans (29)	TAU/waitlist	6	4 (14)	39 (80) overall (including the waitlist that received EFT)	63.23 (PCL completer)	+0.52 (PCL completer)	NR	NR	NR

(continued)

Table 2. Summary of Outcomes From RCTs of Second-Line Interventions for Military-Related PTSD (continued)

Source	Participants (No.)	Conditions	Longest Follow-up, mo	Drop Out	Remission at Post-Treatment	Mean Post-Treatment Total PTSD Score	Pre-Post Drop in PTSD Symptoms	No. (%)		Between-Group ES From Pretreatment to Posttreatment
								% Attaining Clinically Meaningful Change At Posttreatment	% Retaining PTSD Diagnosis at Posttreatment	
Kip et al. ⁴⁷ 2013	Service members and veterans (29)	ART	3	3 (10)	NR	42.00 (PCL ITT)	15.4 (PCL ITT)	17 (65) (PCL completer)	NR	d = 1.25
Engel et al. ⁴⁸ 2014	Service members and veterans (28)	Attention control	3	NR	NR	54.3 (PCL ITT)	2.1 (PCL ITT)	3 (13) (PCL completer)	NR	
Engel et al. ⁴⁸ 2014	Active duty soldiers (28)	Acupuncture + TAU	12 wk	1 (4)	NR	38.8 (PCL ITT)	19.3 (PCL ITT)	NR	NR	NR
	Active duty soldiers (27)	TAU	12 wk	0	NR	51.5 (PCL ITT)	3.9 (PCL ITT)	NR	NR	NR
Kuckertz et al. ⁴⁹ 2014	Inpatient active duty personnel (20)	Attention bias modification	None	8 (40)	NR	42.83 (PCL completer)	20.25 (PCL completer)	NR	NR	NR
	Inpatient active duty personnel (17)	Attention control condition	None	0	NR	51.65 (PCL completer)	10.06 (PCL completer)	NR	NR	NR
Moradi et al. ⁵⁰ 2014	Iranian veterans (12)	MEST	3	0	NR	50.17 (IES-R ITT)	30.91 (IES-R ITT)	NR	NR	d = 4.79
	Iranian veterans (12)	Waitlist	3	0	NR	75.34 (IES-R ITT)	+ 0.01 (IES-R ITT)	NR	NR	

Abbreviations: ART, accelerated resolution therapy; BA-TE, behavioral activation and therapeutic exposure; CAPS, Clinician Administered PTSD Scale; CBT, cognitive behavioral therapy; completer, treatment completer sample only; EFT, emotional freedom techniques; ES, effect size; HT + GI, healing touch with guided imagery; IES-R, Impact of Event Scale-Revised; ITT, intent-to-treat; MBSR, mindfulness-based stress reduction; MEST, memory specificity training; η^2_p , partial eta-squared; NR, not reported; PCL, PTSD Checklist; PCT, present-centered therapy; PSSI, PTSD Symptom Scale Interview; PTSD, posttraumatic stress disorder; TAU, treatment as usual.

1 notable exception,²⁴ sample sizes have tended to be smaller, and no large methodologically robust trials of prolonged exposure have been published in US male veterans with combat trauma. The only available data on prolonged exposure in US combat-exposed male veterans come from 2 small trials that studied biomarkers associated with clinical outcomes.^{26,27}

Efficacy of EMDR

RCTs of EMDR for military-related PTSD (all conducted prior to the wars in Iraq and Afghanistan) involved small samples, tested only brief interventions (1-3 sessions),^{28,55,56} or involved dismantling comparisons (eliminating eye movements).⁵⁷ They also generally did not use methodology consistent with modern trials. In the 2 trials testing adequate doses of EMDR there were large symptom reductions,⁵⁸ and 78% of completers no longer met criteria for PTSD.²⁹ EMDR performed comparably to variants in which the eye tracking was removed or when compared with an active non-trauma-focused therapy.⁵⁸ EMDR outperformed waitlisting (ie, no treatment) and biofeedback-assisted relaxation, with results maintained at 9-month follow-up.²⁹ In sum, the efficacy of EMDR remains largely based on civilian studies⁵⁹; additional studies in military populations are needed.

Second-Line Interventions

Given high dropout and nonresponse rates from first-line therapies, an increasing number of trials have examined an array of alternatives, including variations of cognitive-behavioral therapy^{38,44} or novel delivery modalities, such as virtual reality^{36,39} or web-based content.³³ A variant of EMDR, accelerated resolution therapy, demonstrated large effects ($d = 1.25$) compared with an educational control condition similar to waitlisting in a preliminary RCT of veterans with PTSD.⁴⁷

Other trials have tested complementary and alternative therapies that are theoretically and mechanistically distinct. Acupuncture combined with usual care outperformed usual care alone ($d = 1.7$)⁴⁸; small trials have shown some evidence of efficacy for mindfulness-based interventions,⁴³ mantram repetition (ie, silent repetition of a word or phrase with spiritual significance),^{34,45} attention bias modification,⁴⁹ and memory specificity training.⁵⁰ Healing touch therapy, involving tapping body points while engaging in non-exposure-based guided imagery, also demonstrated efficacy over treatment as usual ($d = 0.85$).⁴¹ These findings suggest that a variety of disparate treatment mechanisms are associated with reduced PTSD symptoms.

In contrast, treatments that have failed to demonstrate efficacy among veterans include relaxation, deep breathing, biofeedback, and mindfulness-based stress reduction.^{30,35,42} A brief 3-session intervention consisting of writing about combat traumas produced little meaningful PTSD improvement and did not outperform writing about one's use of time.⁴⁰ Cognitive-behavioral group therapy aimed at comorbid PTSD and depression,³¹ and

comorbid PTSD and substance dependence, did not substantially improve symptoms and was equivalent to controls.

Discussion

The past 10 years has seen unprecedented interest in identifying effective PTSD treatments for service members and veterans. Findings from RCTs indicate a significant need for further treatment development and improvement. CPT and prolonged exposure, the 2 most widely used first-line (ie, recommended) therapies, show large within-group (pretreatment to posttreatment) effect sizes. However, effect sizes, which are more commonly used in psychology literature than in medical literature, reflect mean outcomes and do not adequately capture heterogeneity in patient outcomes; between one-third and one-half of patients receiving CPT or prolonged exposure did not demonstrate clinically meaningful symptom change (when this outcome was reported). Approximately two-thirds of patients receiving CPT or prolonged exposure retained their diagnosis posttreatment. Mean PTSD scores have tended to remain at or above diagnostic thresholds after treatment, and the 2 studies reporting remission rates suggest that symptom remission is relatively uncommon. Most importantly, many trials of CPT and prolonged exposure have compared patients receiving the intervention with patients not receiving any standardized intervention (waitlist) or with patients receiving treatment as usual. When CPT and prolonged exposure were compared with non-trauma-focused psychotherapies, such as present-centered therapy, similar levels of symptom improvement were often observed, particularly at follow-up intervals.

Approximately one-fourth of patients enrolled in clinical trials and receiving CPT and prolonged exposure dropped out during treatment. These proportions are broadly comparable to the proportions of dropouts in studies of trauma-focused therapies in civilians⁶⁰ and trials of depression in veterans.⁶¹ Treatment nonretention has been a significant problem in military-related PTSD care more generally; several large observational studies in both the VA and DoD found that only a small proportion of individuals receive a minimally adequate number of mental health encounters after PTSD diagnosis.⁶² Reasons for not seeking treatment and dropout are complex and include stigma, confidentiality concerns, time demands, perceived treatment inefficacy, and discomfort with the therapist.⁶²

Current VA policies emphasize CPT and prolonged exposure as treatments of choice. Clinical practice guidelines (including the VA/DoD guideline⁵²), based largely on studies in civilians, also include CPT and prolonged exposure as first-line recommendations for adults with PTSD, although EMDR and other trauma-focused therapies are given at least equal standing (separate recommendations for service members and veterans are not made). Some evidence suggests that outcomes for PTSD treatment tend to be better among civilians than among veterans,^{63,64} although this has not been consistent and remains an empirical question. Potential reasons why treatment outcomes may be worse among military and veteran populations include the extended, repeated, and intense nature of deployment trauma⁶⁵ and the fact that service members are exposed not only to life threats but to traumatic losses and morally compromising experiences that may require different treatment approaches.⁶⁶⁻⁶⁸ A recent meta-analysis comparing trauma-

Box. Descriptions of First-Line Interventions

Cognitive therapy. Focuses on modifying dysfunctional thoughts, beliefs, and expectations by identifying, challenging, and replacing maladaptive cognitions. Cognitive processing therapy (CPT) is the most widely used example of cognitive therapy in the Departments of Defense and Veterans Affairs.

Exposure therapy. Comprises psychoeducation, imaginal or narrative exposure (targeting trauma memories), in vivo exposure (targeting external stimuli or situations that the patient avoids because of the trauma), and processing of thoughts and emotions, with the aim of confronting, rather than avoiding, feared memories and stimuli. Prolonged exposure therapy (PE) is the most widely used example of exposure therapy in the Departments of Defense and Veterans Affairs.

Eye movement desensitization and reprocessing (EMDR). Asks patients to attend to emotionally disturbing material in brief sequential doses while focusing on an external stimulus, typically therapist-directed lateral eye movements. Additionally, treatment involves identifying bodily sensations associated with the image, identifying an aversive cognition associated with the trauma, and identifying an alternative positive cognition to replace the aversive cognition.

Stress inoculation training (SIT). Teaches anxiety-management skills including relaxation training, breathing retraining, positive thinking and self-talk, assertiveness training, and thought stopping. It may also include cognitive restructuring and exposure, although these are optional elements.

focused and non-trauma-focused therapies (both civilian and military trauma) found that, in populations with more complex trauma, such as veterans and refugees, there was little difference in efficacy; moderate differences favoring trauma-focused therapies were only present for less complex traumas.⁶⁹ Additional likely reasons for worse outcomes in veterans include comorbidities (eg, 87% of veterans with PTSD presenting to VA primary care clinics have at least 1 psychiatric comorbidity, with the mode being 3-4 disorders⁷⁰) and disability compensation incentives. To contextualize these findings, a recent narrative review of cognitive-behavioral therapy for depression in veterans similarly showed that relatively few trials are available, cognitive-behavioral therapy often does not outperform controls, and results compare unfavorably with civilian outcomes.⁶¹

One unresolved issue is whether focusing on the trauma, either through exposure or cognitive reframing, is necessary for recovery. The findings that interventions such as present-centered therapy and, in civilians, interpersonal psychotherapy⁷¹ are associated with efficacy similar to that of trauma-focused therapy needs to be reconciled with the common assumption in the field that fidelity to a trauma-focused approach is essential for symptom improvement. VA/DoD treatment guidelines, and other guidelines, specify that patient preference should be a guiding factor in treatment selection. Yet little research has been conducted on patient preferences or on other behavioral and biological predictors of dropping out of care, clearly the strongest influence on treatment effectiveness.⁶²

Several large-scale military-related PTSD trials are currently ongoing, including trials with active-duty service members, a population rarely studied but an important target for early interventions.

A large (N = 900) multisite trial is directly comparing CPT and prolonged exposure in veterans,⁷² although it remains to be seen to what extent continued focus on efficacy comparisons will improve care, particularly since equivalency between leading interventions has consistently been demonstrated in civilians.^{59,73} Ongoing trials focus almost exclusively on 2 trauma-focused interventions, despite these therapies being rarely used in actual VA and DoD clinical practice (less than 10% of the time, in one estimate of New England region VA centers⁷⁴). The effectiveness of psychotherapies for PTSD as actually delivered in the VA and DoD has received little empirical attention.⁷⁵

Limitations of this descriptive review (which did not use formal meta-analytic techniques or access all international databases) include the relatively small number of studies and sample sizes, limiting generalizability. Studies often did not report key outcomes, such as symptom remission, loss of diagnosis, or clinically important subthreshold PTSD^{76,77} (which would include individuals who met previous diagnostic criteria but no longer meet the latest criteria).⁷⁸ It is also noteworthy that the metrics of meaningful symptom improvement (typically a 10- or 12-point decrease in PTSD scores) are researcher-defined, not patient-defined; patient perspectives and preferences have not been a primary focus of research. Likewise, most trials have reported only total PTSD symptom scores, and the potential for differential treatment effects across symptom clusters remains unexamined. Trials also have not reported the need for continued care following CPT or prolonged exposure; for many patients, 12 sessions of manualized trauma- or non-trauma-focused treatment is insufficient. Definitions of treatment dropout also differ between studies, and studies often fail to delineate why patients dropped out. Moreover, military personnel and veterans participating in PTSD trials are often taking psychotropic medications concurrently (approximately three-fourths of participants in CPT and prolonged exposure trials), creating potential confounding. Data are lacking on the relative efficacy of psychotherapy compared with medication or the synergistic effects of combined treatments.⁷⁹ Last, although we considered group and individual therapy together, these 2 modalities are practically and mechanistically distinct. A recent meta-analysis of civilian and veteran group therapies

found smaller effect sizes in combat samples; although group therapy outperformed waitlist controls, it did not outperform active-comparison conditions.⁸⁰

Conclusions

PTSD in military and veteran populations is a complex and difficult-to-treat disorder for which first-line trauma-focused psychotherapy approaches are not optimal. Although efficacious for some patients, first-line treatments have high nonresponse and dropout rates, and patients often remain symptomatic. Two principal clinical conclusions can be drawn from this review. First, the available evidence supports the use of either trauma-focused or structured non-trauma-focused therapies, depending on patient preferences or other factors that might promote treatment retention. Second, there is a need for improvement in existing PTSD treatments as well as the development and testing of novel evidence-based treatment strategies, whether trauma-focused or non-trauma-focused. An increasing number of novel therapeutic approaches have been shown efficacious to varying degrees, but progress in the field is unlikely without better understanding of treatment mechanisms, patient preferences, factors influencing treatment engagement and retention, and behavioral and biomarker prediction of differential treatment responses.

Clinical Bottom Line

- Posttraumatic stress disorder (PTSD) is a disabling psychiatric condition common among military personnel and veterans
- A range of psychotherapies are available, but military-PTSD is complex and difficult to treat
- The available evidence supports the use of structured trauma-focused or non-trauma-focused approaches
- Although trauma-focused and non-trauma-focused interventions often improve symptoms, many patients continue to meet criteria for PTSD after treatment
- There is an urgent need for innovative treatment strategies, whether trauma-focused or non-trauma-focused

ARTICLE INFORMATION

Author Contributions: Dr Steenkamp had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Steenkamp, Litz.

Acquisition, analysis, or interpretation of data: Steenkamp, Litz, Hoge, Marmar.

Drafting of the manuscript: Steenkamp, Litz, Marmar.

Critical revision of the manuscript for important intellectual content: Hoge, Marmar.

Administrative, technical, or material support: Litz. **Study supervision:** Litz, Marmar.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

Disclaimer: The views of this article are those of the authors and do not represent an official

position of the US Army, Department of Defense, or listed institutions.

Additional Contributions: We thank Carol Hundert, BS (VA Boston Healthcare System), and Amelia Tankersley, BA (VA Boston Healthcare System), for assistance with manuscript preparation. They received no compensation for their contributions.

Submissions: We encourage authors to submit papers for consideration as a Review. Please contact Mary McGrae McDermott, MD, at mdm608@northwestern.edu.

REFERENCES

1. Kok BC, Herrell RK, Thomas JL, Hoge CW. Posttraumatic stress disorder associated with combat service in Iraq or Afghanistan: reconciling prevalence differences between studies. *J Nerv Ment Dis.* 2012;200(5):444-450.

2. Kang HK, Natelson BH, Mahan CM, Lee KY, Murphy FM. Post-traumatic stress disorder and chronic fatigue syndrome-like illness among Gulf War veterans: a population-based survey of 30,000 veterans. *Am J Epidemiol.* 2003;157(2):141-148.

3. Marmar CR, Schlenger W, Henn-Haase C, et al. Course of posttraumatic stress disorder 40 years after the Vietnam War: findings from the National Vietnam Veterans Longitudinal Study [published online July 22, 2015]. *JAMA Psychiatry.* doi:10.1001/jamapsychiatry.2015.0803.

4. Kulka RA, Schlenger WE, Fairbank JA, et al. *Trauma and the Vietnam War Generation: Report of Findings From the National Vietnam Veterans Readjustment Study.* New York, NY: Brunner/Mazel; 1990.

5. Hoge CW, Terhakopian A, Castro CA, Messer SC, Engel CC. Association of posttraumatic stress disorder with somatic symptoms, health care visits,

- and absenteeism among Iraq war veterans. *Am J Psychiatry*. 2007;164(1):150-153.
6. Savoca E, Rosenheck R. The civilian labor market experiences of Vietnam-era veterans: the influence of psychiatric disorders. *J Ment Health Policy Econ*. 2000;3(4):199-207.
 7. Riggs DS, Byrne CA, Weathers FW, Litz BT. The quality of the intimate relationships of male Vietnam veterans: problems associated with posttraumatic stress disorder. *J Trauma Stress*. 1998;11(1):87-101.
 8. McFall M, Fontana A, Raskind M, Rosenheck R. Analysis of violent behavior in Vietnam combat veteran psychiatric inpatients with posttraumatic stress disorder. *J Trauma Stress*. 1999;12(3):501-517.
 9. Buckley TC, Mozley SL, Bedard MA, Dewulf AC, Greif J. Preventive health behaviors, health-risk behaviors, physical morbidity, and health-related role functioning impairment in veterans with post-traumatic stress disorder. *Mil Med*. 2004;169(7):536-540.
 10. Zatzick DF, Marmar CR, Weiss DS, et al. Posttraumatic stress disorder and functioning and quality of life outcomes in a nationally representative sample of male Vietnam veterans. *Am J Psychiatry*. 1997;154(12):1690-1695.
 11. Xue C, Ge Y, Tang B, et al. A meta-analysis of risk factors for combat-related PTSD among military personnel and veterans. *PLoS One*. 2015;10(3):e0120270.
 12. Tennant C, Goulston K, Dent O. Clinical psychiatric illness in prisoners of war of the Japanese: forty years after release. *Psychol Med*. 1986;16(4):833-839.
 13. Karlin BE, Ruzek JI, Chard KM, et al. Dissemination of evidence-based psychological treatments for posttraumatic stress disorder in the Veterans Health Administration. *J Trauma Stress*. 2010;23(6):663-673.
 14. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed, text revision. Washington, DC: American Psychiatric Association; 2000.
 15. Monson CM, Schnurr PP, Resick PA, Friedman MJ, Young-Xu Y, Stevens SP. Cognitive processing therapy for veterans with military-related posttraumatic stress disorder. *J Consult Clin Psychol*. 2006;74(5):898-907.
 16. Forbes D, Lloyd D, Nixon RDV, et al. A multisite randomized controlled effectiveness trial of cognitive processing therapy for military-related posttraumatic stress disorder. *J Anxiety Disord*. 2012;26(3):442-452.
 17. Surís A, Link-Malcolm J, Chard K, Ahn C, North C. A randomized clinical trial of cognitive processing therapy for veterans with PTSD related to military sexual trauma. *J Trauma Stress*. 2013;26(1):28-37.
 18. Morland LA, Mackintosh MA, Greene CJ, et al. Cognitive processing therapy for posttraumatic stress disorder delivered to rural veterans via telemental health: a randomized noninferiority clinical trial. *J Clin Psychiatry*. 2014;75(5):470-476.
 19. Resick PA, Wachen JS, Mintz J, et al. A randomized clinical trial of group cognitive processing therapy compared with group present-centered therapy for PTSD among active duty military personnel [published online May 4, 2015]. *J Consult Clin Psychol*. doi:10.1037/ccp0000016.
 20. Cooper NA, Clum GA. Imaginal flooding as a supplementary treatment for PTSD in combat veterans: a controlled study. *Behav Ther*. 1989;20(3):381-391. doi:10.1016/S0005-7894(89)80057-7.
 21. Keane TM, Fairbank JA, Caddell JM, Zimering RT. Implosive (flooding) therapy reduces symptoms of PTSD in Vietnam combat veterans. *Behav Ther*. 1989;20(2):245-260. doi:10.1016/S0005-7894(89)80072-3.
 22. Boudewyns PA, Hyer LA. Physiological response to combat memories and preliminary treatment outcome in Vietnam veteran PTSD patients treated with direct therapeutic exposure. *Behav Ther*. 1990;21(1):63-87. doi:10.1016/S0005-7894(05)80189-3.
 23. Schnurr PP, Friedman MJ, Foy DW, et al. Randomized trial of trauma-focused group therapy for posttraumatic stress disorder: results from a department of veterans affairs cooperative study. *Arch Gen Psychiatry*. 2003;60(5):481-489.
 24. Schnurr PP, Friedman MJ, Engel CC, et al. Cognitive behavioral therapy for posttraumatic stress disorder in women: a randomized controlled trial. *JAMA*. 2007;297(8):820-830.
 25. Nacasch N, Foa EB, Huppert JD, et al. Prolonged exposure therapy for combat- and terror-related posttraumatic stress disorder: a randomized control comparison with treatment as usual. *J Clin Psychiatry*. 2011;72(9):1174-1180.
 26. Yehuda R, Pratchett LC, Elmes MW, et al. Glucocorticoid-related predictors and correlates of post-traumatic stress disorder treatment response in combat veterans. *Interface Focus*. 2014;4(5):20140048.
 27. Rauch SA, King AP, Abelson J, et al. Biological and symptom changes in posttraumatic stress disorder treatment: a randomized clinical trial. *Depress Anxiety*. 2015;32(3):204-212.
 28. Rogers S, Silver SM, Goss J, Obenchain J, Willis A, Whitney RL. A single session, group study of exposure and Eye Movement Desensitization and Reprocessing in treating Posttraumatic Stress Disorder among Vietnam War veterans: preliminary data. *J Anxiety Disord*. 1999;13(1-2):119-130.
 29. Carlson JG, Chemtob CM, Rusnak K, Hedlund NL, Muraoka MY. Eye movement desensitization and reprocessing (EMDR) treatment for combat-related posttraumatic stress disorder. *J Trauma Stress*. 1998;11(1):3-24.
 30. Watson CG, Tuorila JR, Vickers KS, Gearhart LP, Mendez CM. The efficacies of three relaxation regimens in the treatment of PTSD in Vietnam War veterans. *J Clin Psychol*. 1997;53(8):917-923.
 31. Dunn NJ, Rehm LP, Schillaci J, et al. A randomized trial of self-management and psychoeducational group therapies for comorbid chronic posttraumatic stress disorder and depressive disorder. *J Trauma Stress*. 2007;20(3):221-237.
 32. Frueh BC, Monnier J, Yim E, Grubaugh AL, Hamner MB, Knapp RG. A randomized trial of telepsychiatry for post-traumatic stress disorder. *J Telemed Telecare*. 2007;13(3):142-147.
 33. Litz BT, Engel CC, Bryant RA, Papa A. A randomized, controlled proof-of-concept trial of an Internet-based, therapist-assisted self-management treatment for posttraumatic stress disorder. *Am J Psychiatry*. 2007;164(11):1676-1683.
 34. Bormann JE, Thorp S, Wetherell JL, Golshan S. A spiritually based group intervention for combat veterans with posttraumatic stress disorder: feasibility study. *J Holist Nurs*. 2008;26(2):109-116.
 35. Lande RG, Williams LB, Francis JL, Gragnani C, Morin ML. Efficacy of biofeedback for post-traumatic stress disorder. *Complement Ther Med*. 2010;18(6):256-259.
 36. Ready DJ, Gerardi RJ, Backscheider AG, Mascaro N, Rothbaum BO. Comparing virtual reality exposure therapy to present-centered therapy with 11 U.S. Vietnam veterans with PTSD. *Cyberpsychol Behav Soc Netw*. 2010;13(1):49-54.
 37. Beidel DC, Frueh BC, Uhde TW, Wong N, Mentrakoski JM. Multicomponent behavioral treatment for chronic combat-related posttraumatic stress disorder: a randomized controlled trial. *J Anxiety Disord*. 2011;25(2):224-231.
 38. Kent M, Davis MC, Stark SL, Stewart LA. A resilience-oriented treatment for posttraumatic stress disorder: results of a preliminary randomized clinical trial. *J Trauma Stress*. 2011;24(5):591-595.
 39. McLay RN, Wood DP, Webb-Murphy JA, et al. A randomized, controlled trial of virtual reality-graded exposure therapy for post-traumatic stress disorder in active duty service members with combat-related post-traumatic stress disorder. *Cyberpsychol Behav Soc Netw*. 2011;14(4):223-229.
 40. Possemato K, Ouimette P, Kwonlton P. A brief self-guided telehealth intervention for post-traumatic stress disorder in combat veterans: a pilot study. *J Telemed Telecare*. 2011;17(5):245-250.
 41. Jain S, McMahon GF, Hasen P, et al. Healing Touch with Guided Imagery for PTSD in returning active duty military: a randomized controlled trial. *Mil Med*. 2012;177(9):1015-1021.
 42. Kearney DJ, McDermott K, Malte C, Martinez M, Simpson TL. Effects of participation in a mindfulness program for veterans with posttraumatic stress disorder: a randomized controlled pilot study. *J Clin Psychol*. 2013;69(1):14-27.
 43. Niles BL, Klunk-Gillis J, Ryngala DJ, Silberbogen AK, Paysnick A, Wolf EJ. Comparing mindfulness and psychoeducation treatments for combat-related PTSD using a telehealth approach. *Psychol Trauma*. 2012;4(5):538-547. doi:10.1037/a0026161.
 44. Strachan M, Gros DF, Ruggiero KJ, Lejuez CW, Acierno R. An integrated approach to delivering exposure-based treatment for symptoms of PTSD and depression in OIF/OEF veterans: preliminary findings. *Behav Ther*. 2012;43(3):560-569.
 45. Bormann JE, Thorp SR, Wetherell JL, Golshan S, Lang AJ. Meditation-based mantram intervention for veterans with posttraumatic stress disorder: a randomized trial. *Psychol Trauma*. 2013;5(3):259-267. doi:10.1037/a0027522.
 46. Church D, Hawk C, Brooks AJ, et al. Psychological trauma symptom improvement in veterans using emotional freedom techniques: a randomized controlled trial. *J Nerv Ment Dis*. 2013;201(2):153-160.
 47. Kip KE, Rosenzweig L, Hernandez DF, et al. Randomized controlled trial of accelerated resolution therapy (ART) for symptoms of combat-related post-traumatic stress disorder (PTSD). *Mil Med*. 2013;178(12):1298-1309.

48. Engel CC, Cordova EH, Benedek DM, et al. Randomized effectiveness trial of a brief course of acupuncture for posttraumatic stress disorder. *Med Care*. 2014;52(12)(suppl 5):S57-S64.
49. Kuckertz JM, Amir N, Boffa JW, et al. The effectiveness of an attention bias modification program as an adjunctive treatment for post-traumatic stress disorder. *Behav Res Ther*. 2014;63:25-35.
50. Moradi AR, Moshirpanahi S, Parhon H, Mirzaei J, Dalgleish T, Jobson L. A pilot randomized controlled trial investigating the efficacy of Memory Specificity Training in improving symptoms of posttraumatic stress disorder. *Behav Res Ther*. 2014;56:68-74.
51. Forbes D, Creamer M, Bisson JI, et al. A guide to guidelines for the treatment of PTSD and related conditions. *J Trauma Stress*. 2010;23(5):537-552.
52. US Department of Veterans Affairs (VA) and Department of Defense. VA/DoD Clinical Practice Guideline for Management of Post-Traumatic Stress. VA website. <http://www.healthquality.va.gov/guidelines/MH/ptsd/cpgPTSDFULL201011612c.pdf>. 2010. Accessed July 6, 2015.
53. Karlin BE, Cross G. From the laboratory to the therapy room: national dissemination and implementation of evidence-based psychotherapies in the U.S. Department of Veterans Affairs Health Care System. *Am Psychol*. 2014;69(1):19-33.
54. Ruzek JI, Karlin BE, Zeiss A. Implementation of evidence-based psychological treatments in the Veterans Health Administration. In: McHugh RK, Barlow DH, eds. *Dissemination and Implementation of Evidence-Based Psychological Interventions*. New York, NY: Oxford University Press; 2012:78-96.
55. Devilly GJ, Spence SH, Rapee RM. Statistical and reliable change with eye movement desensitization and reprocessing: treating trauma within a veteran population. *Behav Ther*. 1998;29(3):435-455. doi:10.1016/S0005-7894(98)80042-7.
56. Jensen JA. An investigation of eye movement desensitization and reprocessing (EMD/R) as a treatment for posttraumatic stress disorder (PTSD) symptoms of Vietnam combat veterans. *Behav Ther*. 1994;25(2):311-325. doi:10.1016/S0005-7894(05)80290-4.
57. Pitman RK, Orr SP, Altman B, Longpre RE, Poiré RE, Macklin ML. Emotional processing during eye movement desensitization and reprocessing therapy of Vietnam veterans with chronic posttraumatic stress disorder. *Compr Psychiatry*. 1996;37(6):419-429.
58. Boudewyns PA, Hyer LA. Eye movement desensitization and reprocessing (EMDR) as treatment for post-traumatic stress disorder (PTSD). *Clin Psychol Psychother*. 1996;3(3):185-195.
59. Powers MB, Halpern JM, Ferenschak MP, Gillihan SJ, Foa EB. A meta-analytic review of prolonged exposure for posttraumatic stress disorder. *Clin Psychol Rev*. 2010;30(6):635-641.
60. Schottenbauer MA, Glass CR, Arnkoff DB, Tendick V, Gray SH. Nonresponse and dropout rates in outcome studies on PTSD: review and methodological considerations. *Psychiatry*. 2008;71(2):134-168.
61. Hundt NE, Barrera TL, Robinson A, Cully JA. A systematic review of cognitive behavioral therapy for depression in Veterans. *Mil Med*. 2014;179(9):942-949.
62. Hoge CW, Grossman SH, Auchterlonie JL, Riviere LA, Milliken CS, Wilk JE. PTSD treatment for soldiers after combat deployment: low utilization of mental health care and reasons for dropout. *Psychiatr Serv*. 2014;65(8):997-1004.
63. Bradley R, Greene J, Russ E, Dutra L, Westen D. A multidimensional meta-analysis of psychotherapy for PTSD. *Am J Psychiatry*. 2005;162(2):214-227.
64. Watts BV, Schnurr PP, Mayo L, Young-Xu Y, Weeks WB, Friedman MJ. Meta-analysis of the efficacy of treatments for posttraumatic stress disorder. *J Clin Psychiatry*. 2013;74(6):e541-e550.
65. Price M, Gros DF, Strachan M, Ruggiero KJ, Acierno R. Combat experiences, pre-deployment training, and outcome of exposure therapy for post-traumatic stress disorder in Operation Enduring Freedom/Operation Iraqi Freedom veterans. *Clin Psychol Psychother*. 2013;20(4):277-285.
66. Stein NR, Mills MA, Arditte K, et al; STRONG STAR Consortium. A scheme for categorizing traumatic military events. *Behav Modif*. 2012;36(6):787-807.
67. Litz BT, Stein N, Delaney E, et al. Moral injury and moral repair in war veterans: a preliminary model and intervention strategy. *Clin Psychol Rev*. 2009;29(8):695-706.
68. Steenkamp MM, Litz BT, Gray MJ, et al. A brief exposure-based intervention for service members with PTSD. *Cognit Behav Pract*. 2011;18(1):98-107. doi:10.1016/j.cbpra.2009.08.006.
69. Gerger H, Munder T, Barth J. Specific and nonspecific psychological interventions for PTSD symptoms: a meta-analysis with problem complexity as a moderator. *J Clin Psychol*. 2014;70(7):601-615.
70. Magruder KM, Frueh BC, Knapp RG, et al. Prevalence of posttraumatic stress disorder in Veterans Affairs primary care clinics. *Gen Hosp Psychiatry*. 2005;27(3):169-179.
71. Markowitz JC, Petkova E, Neria Y, et al. Is exposure necessary? a randomized clinical trial of interpersonal psychotherapy for PTSD. *Am J Psychiatry*. 2015;172(5):430-440.
72. Schnurr PP, Chard KM, Ruzek JI, et al. Design of VA Cooperative Study #591: CERV-PTSD, comparative effectiveness research in veterans with PTSD. *Contemp Clin Trials*. 2015;41:75-84.
73. Benish SG, Imel ZE, Wampold BE. The relative efficacy of bona fide psychotherapies for treating post-traumatic stress disorder: a meta-analysis of direct comparisons. *Clin Psychol Rev*. 2008;28(5):746-758.
74. Shiner B, D'Avolio LW, Nguyen TM, et al. Measuring use of evidence based psychotherapy for posttraumatic stress disorder. *Adm Policy Ment Health*. 2013;40(4):311-318.
75. Institute of Medicine (IOM). Treatment for Posttraumatic Stress Disorder in Military and Veteran Populations: Final Assessment. IOM website. [http://iom.nationalacademies.org/-/media/Files/Report Files/2014/PTSD-II/PTSD-II-RB.pdf](http://iom.nationalacademies.org/-/media/Files/Report%20Files/2014/PTSD-II/PTSD-II-RB.pdf). 2014. Accessed July 6, 2015.
76. Marshall RD, Olfson M, Hellman F, Blanco C, Guardino M, Struening EL. Comorbidity, impairment, and suicidality in subthreshold PTSD. *Am J Psychiatry*. 2001;158(9):1467-1473.
77. Stein MB, Walker JR, Hazen AL, Forde DR. Full and partial posttraumatic stress disorder: findings from a community survey. *Am J Psychiatry*. 1997;154(8):1114-1119.
78. Hoge CW, Riviere LA, Wilk JE, Herrell RK, Weathers FW. The prevalence of post-traumatic stress disorder (PTSD) in US combat soldiers: a head-to-head comparison of DSM-5 vs DSM-IV-TR symptom criteria with the PTSD checklist. *Lancet Psychiatry*. 2014;1(4):269-277. doi:10.1016/S2215-0366(14)70235-4.
79. Rothbaum BO, Price M, Jovanovic T, et al. A randomized, double-blind evaluation of D-cycloserine or alprazolam combined with virtual reality exposure therapy for posttraumatic stress disorder in Iraq and Afghanistan War veterans. *Am J Psychiatry*. 2014;171(6):640-648.
80. Sloan DM, Feinstein BA, Gallagher MW, Beck JG, Keane TM. Efficacy of group treatment for posttraumatic stress disorder symptoms: a meta-analysis. *Psychol Trauma*. 2013;5(2):176-183. doi:10.1037/a0026291.