GRAIL Baseline Assessment		
	Administered Assessment	Description
1.	Demographics	Participant demographics and socio-economic status
2.	HIV Testing	Asks when they first test positive for HIV
3.	ART Medication History and Adherence	History and current use of ART, VAS for adherence
4.	Resource Utilization	Assesses hospital stays
5.	Alcohol Use Disorders Identification Test (AUDIT)	Identifies persons with hazardous alcohol use during the past year
6.	DSM-5 Alcohol Use Disorder	Diagnosis of alcohol use disorders, with classifications of mild, moderate, and severe
7.	30-day Alcohol Timeline Followback (TLFB)	Estimates of participant's daily drinking over the past 30 days
8.	Penn Alcohol Craving Scale	Instrument for assessing alcohol craving. Measures frequency, intensity, and duration of thoughts about drinking, as well as ability to resist drinking
9.	Readiness to Quit Alcohol	Assesses readiness to quit drinking via a readiness to change ruler
10.	Tobacco Use (Fagerström)	A measure of nicotine dependence
11.	HIV Risk Categories	Asks about participant's potential exposure to HIV
12.	Depressive Symptoms (CESD-10) ⁸	A measure of depressive symptomology
13.	Anxiety (GAD-7)	Measure of severity of anxiety
14.	Co-Morbidities	Participant and family medical history
15.	Brief Pain Inventory (SF)	Assesses the severity of pain and its impact on functioning
16.	Medications	Prescription and non-prescription medication taken in the past 7 days
17.	Drug Use	Participant's self-reported drug use, modified to adapt to drug practices in Uganda and questions regarding history of overdose
18.	Social Support Scale	Measures access to companionship, assistance, or other types of support
19.	Veterans Rand 12-Item Health Survey	Assessment of overall health
20.	Symptoms/events Checklist	Assesses baseline and follow-up symptoms