

1 RESEARCH ARTICLE

2 RUNNING HEAD: Eye closure impacts visual processing

3 Differential cortical and subcortical visual processing with  
4 eyes shut

5 Nicholas G. Cicero,<sup>1,3,4</sup> Michaela Klimova,<sup>2</sup> Laura D. Lewis\*,<sup>3,4,5,6</sup> Sam Ling\*<sup>1,2</sup>

6 <sup>1</sup>Graduate Program in Neuroscience, Boston University, Boston, MA

7 <sup>2</sup>Department of Psychological and Brain Sciences, Boston University, Boston, MA

8 <sup>3</sup>Department of Biomedical Engineering, Boston University, Boston, MA

9 <sup>4</sup>Institute for Medical Engineering and Sciences, Massachusetts Institute of Technology, Cambridge, MA

10 <sup>5</sup>Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology,  
11 Cambridge, MA

12 <sup>6</sup>Athinoula A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General  
13 Hospital, Charlestown, MA

14

15 Correspondence: Sam Ling ([samling@bu.edu](mailto:samling@bu.edu)) and Laura Lewis ([ldlewis@mit.edu](mailto:ldlewis@mit.edu))

16 36-724 50 Vassar St., Massachusetts Institute of Technology, Cambridge, MA 02139

17

---

18 **ABSTRACT**

19 Closing our eyes largely shuts down our ability to see. That said, our eyelids still pass some light,  
20 allowing our visual system to coarsely process information about visual scenes, such as changes in  
21 luminance. However, the specific impact of eye closure on processing within the early visual system  
22 remains largely unknown. To understand how visual processing is modulated when eyes are shut, we  
23 used functional magnetic resonance imaging (fMRI) to measure responses to a flickering visual stimulus  
24 at high (100%) and low (10%) temporal contrasts, while participants viewed the stimuli with their eyes  
25 open or closed. Interestingly, we discovered that eye closure produced a qualitatively distinct pattern of  
26 effects across the visual thalamus and visual cortex. We found that with eyes open, low temporal  
27 contrast stimuli produced smaller responses, across the lateral geniculate nucleus (LGN), primary (V1)  
28 and extrastriate visual cortex (V2). However, with eyes closed, we discovered that the LGN and V1  
29 maintained similar BOLD responses as the eyes open condition, despite the suppressed visual input  
30 through the eyelid. In contrast, V2 and V3 had strongly attenuated BOLD response when eyes were  
31 closed, regardless of temporal contrast. Our findings reveal a qualitatively distinct pattern of visual  
32 processing when the eyes are closed – one that is not simply an overall attenuation, but rather reflects  
33 distinct responses across visual thalamocortical networks, wherein the earliest stages of processing  
34 preserves information about stimuli but is then gated off downstream in visual cortex.

35 **NEW & NOTEWORTHY**

36 When we close our eyes coarse luminance information is still accessible by the visual system. Using  
37 functional magnetic resonance imaging, we examined whether eyelid closure plays a unique role in  
38 visual processing. We discovered that while the LGN and V1 show equivalent responses when the eyes

39 are open or closed, extrastriate cortex exhibited attenuated responses with eye closure. This suggests  
40 that when the eyes are closed, downstream visual processing is blind to this information.

41 **Keywords:** luminance, eye closure, fMRI, LGN, visual cortex

42

---

## 43 INTRODUCTION

44 Light exposure during sleep has substantial effects on the brain: it can alter circadian rhythms,  
45 sleep quality, and mood (1,2). During sleep, our eyes are closed and the eyelids function as potent filters  
46 of visual information. However, our eyelids are only partial filters and do not completely attenuate all  
47 visual information (3,4). The eyelid has been characterized as a red-pass filter, with an estimated 6% red  
48 light spectral transmittance (3). Indeed, subjective experience with high luminance stimuli, such as  
49 during a sunny day, corroborates the idea that changes in luminance are still detectable when our eyes  
50 are closed. With partial, rather than complete, filtering properties, it follows that the visual system  
51 processes external visual information with our eyes closed, as well.

52 How does the visual system process information when our eyes are closed? It is possible that  
53 the filtering properties of the eyelid simply quantitatively suppress responses across visual regions, due  
54 to the attenuation of input. Alternatively, eye closure could induce qualitatively distinct changes in  
55 visual response, selectively modulating responses in specific brain networks. While little is known about  
56 stimulus-evoked visual responses with eyes closed, resting-state fMRI studies have investigated  
57 spontaneous dynamics during eye closure in the absence of any visual stimulus presentation (5-7). These  
58 studies found differences in resting-state functional connectivity in attentional networks depending on  
59 whether eyes were open or closed, along with differences in activation in prefrontal cortex, parietal and  
60 frontal eye fields, and LGN. While eye closure appears to play a unique role in modulating brain  
61 responses, the impact that eye closure has on stimulus-evoked visual responses remains poorly  
62 understood.

63 In this study, we sought to shed light on the role that eye closure plays in modulating responses  
64 within the visual processing hierarchy. To do so, we measured fMRI BOLD responses within visual cortex  
65 and subcortex while participants viewed high and low intensity visual stimuli, with their eyes open or  
66 shut. We manipulated the intensity of visual input via temporal contrast modulation, in which the  
67 luminance of a uniform visual stimuli flickered rapidly between extreme whites and blacks (high  
68 temporal contrast), or between middling intensities (low temporal contrast). Indeed, previous work has  
69 shown visuocortical responses to be sensitive to changes in luminance (8). By measuring BOLD  
70 responses to high and low luminance contrast stimuli, we examined whether there is a qualitatively  
71 unique pattern of luminance responses across the visuocortical hierarchy when one's eyes are closed,  
72 compared to when they are open.

## 73 MATERIALS AND METHODS


### 74 Participants

75 Data was acquired from a total of 8 healthy participants (5 females, 3 males; 3 Asian, 1 Black or  
76 African American; 4 White). Participants were aged 18-35 years, reported normal or corrected-to-  
77 normal visual acuity, and were recruited from Boston University and the surrounding community. All  
78 participants provided written informed consent before study enrollment and completed a metal  
79 screening form indicating that they had no MRI contraindications. Participants were reimbursed for their  
80 study participation.

## 81 Apparatus & stimuli

82 Stimuli were generated using custom software written in MATLAB (version 2019b) in  
83 conjunction with Psychtoolbox (9). Participants viewed stimuli that was back-projected onto a screen set  
84 within the MRI scanner, using a ProPIXX DLP LED (VPixx Technologies) projector system (minimum  
85 luminance: 1.2 cd/m<sup>2</sup>; maximum luminance: 2507.9 cd/m<sup>2</sup>). Photometer measurements (model LS-100;  
86 Konica Minolta) carried out before the study were used to verify the linearity of the display (1 digital-to-  
87 analog conversion (DAC) step = 9.835 cd/m<sup>2</sup>). These measurements were used to calculate the stimulus  
88 luminance and were acquired from the inner-facing side of the back-projection screen while positioned  
89 within the MRI scanner bore. This was done to best account for the attenuation in luminance due to  
90 back-projection screen characteristics.

91 During each functional run, participants fixated on a median luminance crosshair at the center  
92 of the display while shown a full screen flickering display (17 degrees of visual angle) with no spatial  
93 contrast (Figure 1). The full field flicker was presented in a block design with three trial types (baseline,  
94 high, and low temporal luminance contrast), with each event lasting 16 seconds. In the *baseline* events,  
95 the full field display was a constant median luminance with no luminance modulation. During *high*  
96 events, the full field display flickered with an amplitude envelope of 100% around the middle luminance  
97 value. For *low* events the full field display flickered with an amplitude envelope of 10% around the  
98 median luminance value. All high and low events flickered at a frequency of 7 Hz.  
99



100 Figure 1.

101

## 102 Experimental design

103 Subjects participated in two scan sessions, each lasting approximately two hours. The first  
104 session was dedicated to collecting anatomical images and data for population receptive field (pRF)  
105 mapping using standard techniques and stimuli (10,11). The second session was dedicated to collecting  
106 proton-density (PD) weighted anatomical imaging and fMRI blood oxygenation level-dependent (BOLD)  
107 data across the eyes open and closed conditions, during the luminance task.

108 During the second experimental session, we collected three PD-weighted anatomical scans. PD-  
109 weighted anatomical imaging has previously been used to better localize the LGN (12,13). Following the  
110 PD-weighted scans, participants completed three consecutive runs of a functional localizer. The visual  
111 stimulus for the functional localizer contained a full field flickering grating stimulus (diameter = 6.0°)  
112 with a centered circle (diameter = 0.8°). Within the centered circle, letters rapidly appeared one at a  
113 time with a new letter appearing every 200 ms. Participants were instructed to press a button whenever  
114 the letters 'J' and 'K' appeared within the centered circle. During the localizer blocks, the full field  
115 display alternated between a flickering grating stimulus and a full field non-flickering display at median  
116 luminance value. Participants completed 12 total blocks (6 flickering field, 6 non-flickering field) with an

117 extra non-flickering block at the beginning of the run. At the end of each localizer run, participants were  
118 asked to report their wakefulness level.

119 Participants then completed the luminance flicker task. The task began and ended with a  
120 baseline event. High and low temporal contrast conditions were pseudo-randomly ordered, with all high  
121 and low events interleaved with a baseline event. Each run contained 12 events (6 high, 6 low)  
122 interspersed with 12 baseline events, lasting a total of 384 seconds. On each run participants were  
123 instructed to press a button after each full breath cycle (1 inhale, 1 exhale). This button task was chosen  
124 to ensure that participants did not fall asleep and engaged with the task, while not requiring eyes to be  
125 open. For each run, participants were instructed to either keep their eyes open and fixate on the  
126 crosshair or to keep their eyes closed throughout the run. Each scan session began with an eyes-closed  
127 run, and consecutive runs alternated between open and closed conditions. We always began with the  
128 eyes closed condition to ensure we acquired a sufficient number of runs in this condition, where BOLD  
129 modulations may be lower compared to eyes-open runs. To ensure participants kept their eyes closed or  
130 open, real time eye monitoring was carried out using an EyeLink1000, for the duration of each run. On  
131 average, we collected 5 runs with eyes closed and 4 runs with eyes open, for each subject.  
132

### 133 MRI data acquisition

134 All neuroimaging data were acquired using a research-dedicated Siemens Prisma 3T scanner  
135 using a Siemens 64-channel head coil. A whole brain anatomical scan was acquired using a T1-weighted  
136 multi-echo MPRAGE (1 mm isotropic voxels; field of view (FOV) = 192 x 192 x 134 mm, flip angle (FA) =  
137 7.00°, repetition time (TR) = 2200 ms, echo time (TE) = 1.57 ms). Proton density (PD)-weighted  
138 anatomical scans were acquired to localize LGN (0.9mm x 0.9mm x 1.7mm; TR = 2950.0 ms; TE = 15.6  
139 ms; FA = 180°). Functional scans were acquired using T2\*-weighted in-plane simultaneous imaging (2  
140 mm isotropic voxels; FOV = 104 x 104 x 70 mm, FA = 64.00°, TR = 1000 ms, TE = 30 ms, SMS factor = 5,  
141 GRAPPA acceleration = 2).

### 142 Anatomical data analysis

143 T1-weighted anatomical data were analyzed using the standard “recon-all” pipeline provided by  
144 the FreeSurfer neuroimaging analysis package (14), generating cortical surface models, whole brain  
145 segmentations, and cortical parcellations. All PD-weighted scans were aligned to each subject’s  
146 anatomical space and averaged together (using AFNI’s 3dcalc).

147

### 148 Functional data analysis

149 Functional BOLD time-series data were first corrected for echo-planar imaging (EPI) distortions  
150 using a reverse phase-encode method implemented in FSL (15) and were then preprocessed with FS-  
151 FAST using standard motion-correction procedures, slice timing correction, and boundary-based  
152 registration between functional and anatomical spaces (16). To optimize spatial precision of  
153 experimental data, no volumetric spatial smoothing was performed (full-width half-maximum 0 mm). To  
154 achieve precise alignment of experimental data within the session, cross-run within-modality robust  
155 rigid registration was performed, using the middle time point of each run (17). BOLD time-series data  
156 were demeaned and converted to units of percent signal change. Data collected during the separate pRF  
157 mapping scans were analyzed using the analyzePRF toolbox (11). Results from the pRF model were used  
158 to manually draw labels for our regions of interest within visual cortex.  
159

## 160 Statistical analysis

161 The results from the pRF modeling were used to identify region-of-interest (ROI) labels for each  
162 cortical region before analysis. ROI labels included voxels located inside the cortical ribbon for  
163 V1/V2/V3, which were identified using a visual area network label generated using an intrinsic  
164 functional connectivity atlas (18). Results from the pRF modeling were additionally used to select voxels  
165 with visual field eccentricity preferences less than 17 degrees visual angle away from fixation as this was  
166 the measured extent of the screen within the MRI scanner. Cortical voxels with a poor pRF model fit ( $r^2 <$   
167  $0.10$ ) were removed from further analyses. Initial LGN labels were acquired from thalamic segmentation  
168 and parcellation in Free-Surfer for each participant. These initial labels were overlaid with the GLM  
169 results from the functional localizer and the PD-weighted scans, and only intersecting voxels were  
170 chosen for the final LGN labels and further analyses.


171 An event-triggered average was computed for each flickering condition (low and high) per eyelid  
172 condition and ROI. The BOLD time-series for each ROI per run was separated by the low and high trials,  
173 and all trials of a given type were averaged together. Average BOLD magnitude in response to the  
174 stimulus presentation was computed by averaging 4-16 s post-stimulus onset for each trial. Two-way  
175 between-subjects ANOVA were performed to test for any main effects of temporal contrast and eye  
176 closure and any interaction of the two on average BOLD magnitude during stimulus presentation.  
177 Additional event-triggered average analysis was done with eccentricity, in which the time-series for  
178 V1/V2/V3 voxels were first separated into eccentricity bins defined by degree visual angle relative to  
179 fixation. Foveal-tuned voxels were between  $0.01^\circ - 1.5^\circ$ , parafoveal-tuned voxels were between  $1.5^\circ -$   
180  $4.0^\circ$ , and peripheral-tuned voxels were between  $4.0^\circ - 17.0^\circ$ . An additional ANOVA was performed to  
181 test for any main effect of eccentricity on BOLD response during stimulus presentation. Multiple  
182 comparison correction was done using Bonferroni correction of  $\alpha/n$  at a familywise  $\alpha$  of 0.05 where  $n$  is  
183 the number of tests performed.  
184

## 185 RESULTS

186 We first examined how temporal contrast modulated thalamic and visuocortical responses, and  
187 if eye closure impacted these responses. With eyes open, LGN, V1, and V2 showed larger responses to  
188 high temporal contrast stimuli, compared to low temporal contrast stimuli (Figure 2A). Indeed, during  
189 eyes open with high temporal contrast stimuli, all ROIs had significantly elevated BOLD responses [LGN:  
190  $t(7) = 3.27$ ,  $P = 0.006$ ; V1:  $t(7) = 5.16$ ,  $P < 0.0001$ ; V2:  $t(7) = 3.40$ ,  $P = 0.005$ ; V3:  $t(7) = 2.54$ ,  $P = 0.019$ ],  
191 though the significant response in V3 did not survive multiple comparisons correction. When the  
192 participants closed their eyes, however, LGN and V1 maintained their stronger responses to higher  
193 contrast stimuli [LGN:  $F(1,31) = 4.31$ ,  $P = 0.047$ ; V1:  $F(1,31) = 11.05$ ,  $P = 0.002$ ], which did not differ from  
194 their eyes closed conditions [LGN:  $F(1,31) = 0.02$ ,  $P = 0.975$ ; V1:  $F(1,31) = 1.74$ ,  $P = 0.20$ ]. In other words,  
195 while responses in LGN and V1 were significantly modulated by temporal contrast, they were  
196 completely unaffected by eye closure, despite the profound suppression of visual input from the eyelid.

197 Interestingly, while eye closure did not appear to have a major effect on the earliest stages of  
198 visual processing (LGN and V1), we observed a qualitatively distinct pattern within extrastriate cortices  
199 V2 and V3. When the eyes were closed, there was a drastic attenuation of stimulus evoked responses,  
200 regardless of temporal contrast [Main effects of eye closure: V2:  $F(1,31) = 6.45$ ,  $P = 0.017$ ; V3:  $F(1,31) =$   
201  $5.79$ ,  $P = 0.02$ ; Main effect of temporal contrast V2:  $F(1,31) = 2.91$ ,  $P = 0.09$ ; V3:  $F(1,31) = 0.54$ ,  $P = 0.47$ ].  
202 Pairwise comparisons revealed a significant decrease in BOLD response to high temporal contrast stimuli  
203 with eye closure in V2 ( $t(14) = -3.13$ ;  $P = 0.003$ ) and V3 ( $t(14) = -3.09$ ;  $P = 0.003$ ). Overall, these results  
204 indicate that visual processing appears to be qualitatively different with eyes closed compared to when  
205 eyes are open. The BOLD response in LGN and V1 was modulated by temporal contrast but was

206 unaffected by eye closure, whereas eye closure strongly reduced responses in extrastriate cortices V2  
207 and V3.



208 **Figure 2.**

209

210

211

212

213

214

215

216

217

218

219

220

221

222


223

224

225

Along with the heterogeneity in patterns observed across striate and extrastriate regions, it is possible that there exists heterogeneity *within* each region. It has been reported that there is an eccentricity bias of the BOLD response in V1 and V2, when participants viewed center-surround stimuli with no local contrast (19). To test for an eccentricity bias and if eye closure impacts this bias, we separated voxels in V1-V3 by their eccentricity preference, based on pRF estimates (LGN was excluded from this analysis due to being underpowered for pRF analyses). We defined foveally-preferring voxels as those preferring between  $0.01^\circ - 1.5^\circ$  from fixation, parafoveal-preferring voxels were those between  $1.5^\circ - 4.0^\circ$ , and peripheral-preferring voxels were between  $4.0^\circ - 17.0^\circ$ . As low temporal contrast trials elicited no significant activation across visuocortical regions, we did not test for an effect of eccentricity during low temporal contrast trials. We found that the effect of eccentricity was not significant in V1 [ $F(2,47) = 1.16, P = 0.333$ ] (Figure 3), nor in V2 [ $F(2,47) = 0.51, P = 0.606$ ] nor V3 [ $F(2,47) = 0.30, P = 0.744$ ]. No ROIs had any significant interaction between eye closure and eccentricity [V1:  $F(2,47) = 0.26, P = 0.772$ ; V2:  $F(2,47) = 0.94, P = 0.397$ ; V3:  $F(2,47) = 0.26, P = 0.768$ ]. This suggests that across striate and extrastriate cortices there is no eccentricity bias in BOLD responses nor any difference with eye closure. Thus, the impact of high temporal contrast stimuli and eye closure on BOLD appear uniform within each visuocortical area.





226 *Figure 3.*

227 **DISCUSSION**

228 With subjective experience it is clear that we can still perceive visual stimuli with closed eyes,  
 229 but how distinct stages of the visual system supported this filtered visual experience was unknown. In  
 230 this study, we found that eye closure produces a qualitatively distinct pattern of modulatory responses  
 231 within the early visual system: closing one's eyes selectively attenuated luminance processing in  
 232 extrastriate cortex, but not in LGN nor striate cortex.

233 In line with previous literature showing that early visual responses can still occur when the eyes  
 234 are closed (5,20), we demonstrated that with closed eyes, luminance-dependent responses remain  
 235 present in the LGN and V1. However, we found substantial heterogeneity in activation across regions  
 236 when eyes were closed. One hypothesis as to why we observed strongly attenuated BOLD with closed  
 237 eyes in extrastriate cortex, but not the LGN nor striate cortex, is that top-down modulation of  
 238 visuocortical responses is often stronger in extrastriate compared to striate cortex (21-23). It has been  
 239 demonstrated that higher-order sensory regions, such as the frontal eye field (FEF), may account for the  
 240 selective top-down modulation of extrastriate cortical responses (24). Resting-state fMRI studies that  
 241 examined altered functional connectivity between eyes open and closed states found increased  
 242 activation of the FEF during eyes closed relative to eyes open scans (7), lending further support to top-  
 243 down modulation of extrastriate cortex during eyes closed states. Interestingly, one study which  
 244 microstimulated the FEF of monkeys and measured visuocortical responses with fMRI found that FEF  
 245 stimulation modulated extrastriate areas only in the presence of a visual stimulus, indicating that top-  
 246 down modulation of the extrastriate cortices is dependent on bottom-up influence (25). Since our  
 247 paradigm includes a visual stimulus, it is possible that eye closure in the presence of visual stimuli  
 248 attenuates extrastriate cortical responses through both top-down and bottom-up mechanisms.  
 249 Additionally, the eyelid abolishes almost all structure and form-like information, which is necessary to  
 250 elicit responses in extrastriate cortices that prefer higher-level feature selectivity, such as spatial  
 251 contrast, shapes, and contours. However, eyelid closure still passes through luminance information,

252 which is known to activate striate cortex (8). This preservation of luminance information, but  
253 attenuation of higher-level information, may explain the preservation of early visual pathway activation  
254 with weakened extrastriate activation.

255 Visuocortical responses have been shown to depend on luminance modulation, with responses  
256 increasing monotonically with luminance (8). In addition to luminance modulation, luminance response  
257 functions are strongly contrast dependent, with lower spatial contrast drastically decreasing  
258 visuocortical responses to luminance (8). Since the eyelid filters out much visual information, it is likely  
259 that spatial contrast no longer impacts visual responses and that luminance information dominates what  
260 might pass through the eyelid. Additionally, the lower luminance retinal input with eye closure cannot  
261 fully explain our results since LGN and V1 showed no significant change in BOLD activation between  
262 open and closed eye conditions. Since the eyelid is characterized as a red-pass filter (3), it is possible  
263 that early visual pathways preferentially process this red visual content that extrastriate cortex is blind  
264 to; however, to our knowledge no evidence of this exists. Although further work will be needed to  
265 better unpack luminance responses in the early visual system, our results suggest that luminance-based  
266 responses within early visual areas may not always necessitate the existence of spatial contrast in order  
267 to reveal themselves, as previously suggested.

268 There did not appear to be any significant effects of eccentricity on luminance responses,  
269 neither with eyes open nor eyes closed. Previous research found an eccentricity bias in BOLD responses  
270 in V1 and V2 to center-surround stimuli with no local contrast, most strongly at the edge between the  
271 center and surround of the stimuli (19). However, we may not have observed a similar effect simply  
272 because our visual stimulus did not include any spatial contrast, thereby precluding any edge effects  
273 from a center-surround stimulus. This suggests that the luminance-dependent effect of eccentricity  
274 might depend on where in the visual field an edge exists, which is consistent with another study  
275 examining luxotonic responses in the visual cortex using fMRI, which also did not find any effect of  
276 eccentricity on luxotonicity (8).

277 Our stimulus was designed to provide similar features of visual input across conditions, by  
278 presenting a diffuse light with no spatial contrast, but the input was nevertheless not identical when the  
279 eyes were closed due to the additional attenuation from the eyelid. Future experiments could test a  
280 stimulus that mimics the reduced retinal input of the eyelid. This would involve measuring the spectral  
281 transmittance of each participant's eyelid and adjusting the high temporal contrast visual stimulus to  
282 account for the attenuated transmittance. Such a stimulus would necessarily abolish all structure and  
283 form uniformly across the visual scene, which under standard models of center-surround neuronal  
284 receptive field organization would predict no net change in visuocortical responses due to equivalent  
285 stimulation across excitatory and inhibitory components of the visual cortex (26,27). However, recent  
286 findings of luminance modulation within the visual cortex suggests that luminance information alone  
287 can drive the visual system (8). Thus, a stimulus mimicking the eyelid preserves luminance information  
288 and would be expected to still activate the visual system, in accordance with our eyes closed condition  
289 in which high intensity stimuli still activates LGN and V1. If the visual thalamus and visuocortical  
290 responses between a stimulus mimicking the eyelid with open eyes and the high intensity stimulus in  
291 our eyes closed condition did not align, this would suggest that the physical properties of the eyelid are  
292 not sufficient to explain our results.

293 An alternative explanation of our results is that modulation of visual processing during eye  
294 closure may be dependent on brain state, not just the physical barrier of the eyelid. Eye closure likely  
295 induces a change in overall brain state that alters both the processing of visual information and large-  
296 scale functional network processing. Eye closure decreases activity in attentional systems in the occipital  
297 and parietal lobes and increases functional coupling between sensory thalamus and somatosensory  
298 regions (5-7). Even during short eye blinks, there is evidence of increased activity in the default mode  
299 network (DMN) and decreased activity in the dorsal attention network, suggesting attentional



300 disengagement even with short eye closures (28). During longer eye closures, as examined in this study,  
301 we might expect prolonged or even greater magnitude DMN activation, supporting this idea of  
302 attentional disengagement with closed eyes. These differences in spontaneous brain activity across  
303 sensory and attentional systems point to altered brain states with eye closure. Exteroceptive and  
304 interoceptive mental state hypotheses have been formulated where an exteroceptive mental state is  
305 characterized by increased attention and sensory processing of the external environment with eyes  
306 open (5). On the other hand, an interoceptive mental state is characterized by internally-directed  
307 cognition and reduced sensory processing with eyes closed. Many brain states require prolonged  
308 periods of eye closure, such as sleep and meditation, that involve reduced sensory awareness of  
309 external stimuli and enhanced internally-directed attention. Thus, eye closure may modulate visual  
310 processing through attentional or brain-state-dependent mechanisms.

311

## 312 **DATA AVAILABILITY**

313 Source data for this study are openly available at DOI: 10.18112/openneuro.ds005194.v1.1.0

## 314 **IRB STATEMENT**

315 All aspects of the study were approved by Boston University's Institutional Review Board.

## 316 **ACKNOWLEDGEMENTS**

317 This research was carried out at the Boston University Cognitive Neuroimaging Center. This work  
318 involved the use of instrumentation supported by the NSF Major Research Instrumentation grant BCS-  
319 1625552. We acknowledge the University of Minnesota Center for Magnetic Resonance Research for  
320 use of the multiband-EPI pulse sequences. Data was analyzed on a high-performance computing cluster  
321 supported by the ONR grant N00014-17-1-2304. We thank Shruthi Chakrapani and Stephanie McMains  
322 for assistance with data collection, and members of the S.L. laboratory and L.D.L. laboratory for helpful  
323 feedback on the manuscript.

## 324 **GRANTS**

325 National Institutes of Health R01 EY028163 (to SL)  
326 Sloan Fellowship (to LDL)  
327 McKnight Scholar Award (to LDL)  
328 Pew Biomedical Scholar Award (to LDL)  
329 NIH U19- NS128613 (to LDL)

## 330 **DISCLOSURES**

331 The authors declare no competing financial interests

## 332 **DISCLAIMERS**

333 The authors declare no disclaimers

## 334 **AUTHOR CONTRIBUTIONS**

335 Conceived and designed research: NGC, MK, LDL, SL  
336 Performed experiments: NGC, MK  
337 Analyzed data: NGC  
338 Interpreted results of experiments: NGC, MK, LDL, SL  
339 Prepared figures: NGC  
340 Drafted manuscript: NGC  
341 Edited and revised manuscript: NGC, MK, LDL, SL  
342 Approved final version of manuscript: NGC, MK, LDL, SL

343  
344

345

346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391

## REFERENCES

1. **Blume C, Garbazza C, Spitschan M.** Effects of light on human circadian rhythms, sleep and mood *Somnologie* 23: 147–156, 2019.
2. **Ohayon MM, Milesi C.** Artificial outdoor nighttime lights associate with altered sleep behavior in the American general population. *Sleep* 39: 1311–1320, 2016.
3. **Ando K, Kripke DF.** Light Attenuation by the Human Eyelid. 1996
4. **Bierman A, Figueiro MG, Rea MS.** Measuring and predicting eyelid spectral transmittance. *Journal of Biomedical Optics* 16: 067011, 2011.
5. **Marx E, Deutschländer A, Stepha T, Dieterich M, Wiesmann M, Brandt T.** Eyes open and eyes closed as rest conditions: Impact on brain activation patterns. *NeuroImage* 21: 1818–1824, 2003.
6. **Wei J, Chen T, Li C, Liu G, Qiu J, Wei D.** Eyes-open and eyes-closed resting states with opposite brain activity in sensorimotor and occipital regions: Multidimensional evidences from machine learning perspective. *Frontiers in Human Neuroscience* 12, 2018
7. **Weng Y, Liu X, Hu H, Huang H, Zheng S, Chen Q, Song J, Cao B, Wang J, Wang S, Huang R.** Open eyes and closed eyes elicit different temporal properties of brain functional networks. *NeuroImage* 222, 2020.
8. **Vinke LN, Ling S.** Luminance potentiates human visuocortical responses. *J Neurophysiol* 123: 473–483, 2020.
9. **Brainard DH.** The Psychophysics Toolbox. *Spatial Vision* 10:433–436, 1997.
10. **Dumoulin SO, Wandell BA.** Population receptive field estimates in human visual cortex. *NeuroImage* 39: 647-660, 2008.
11. **Kay KN, Winawer J, Mezer A, Wandell BA.** Compressive spatial summation in human visual cortex. *J Neurophysiol* 110: 481–494, 2013.
12. **Fujita N, Tanaka H, Takanashi M, Hirabuki N, Abe K, Yoshimura H, Nakamura H.** Lateral geniculate nucleus: Anatomic and functional identification by use of MR imaging. *American Journal of Neuroradiology* 22:1719-1726, 2001.
13. **Ling S, Pratte MS, Tong F.** Attention alters orientation processing in the human lateral geniculate nucleus. *Nature Neuroscience* 18: 496–498, 2015.
14. **Fischl B.** FreeSurfer. *NeuroImage* 62: 774–781, 2012.
15. **Andersson JLR, Skare S, Ashburner J.** How to correct susceptibility distortions in spin-echo echo-planar images: Application to diffusion tensor imaging. *NeuroImage* 20: 870–888, 2003.
16. **Greve DN, Fischl B.** Accurate and robust brain image alignment using boundary-based registration. *NeuroImage* 48: 63–72, 2009.
17. **Reuter, M, Rosas, HD, Fischl, B.** Highly accurate inverse consistent registration: A robust approach. *NeuroImage*, 53: 1181-1196, 2011.
18. **Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, Roffman JL, Smoller JW, Zöllei L, Polimeni JR, Fischl B, Liu H, Buckner RL.** The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* 106: 1125–1165, 2011.
19. **Cornelissen FW, Wade AR, Vladusich T, Dougherty RF, Wandell BA.** No functional magnetic resonance imaging evidence for brightness and color filling-in in early human visual cortex. *Journal of Neuroscience* 26: 3634-3641, 2006.
20. **Sharon O, Nir Y.** Attenuated fast steady-state visual evoked potentials during human sleep. *Cerebral Cortex* 28: 1297-1311, 2018.
21. **Haenny PE, Schiller PH.** State dependent activity in monkey visual cortex. *Experimental Brain Research* 69: 225–244, 1988.

- 392 22. Moran J, Desimone R. Selective Attention Gates Visual Processing in the Extrastriate Cortex.  
393 *Science* 229: 782–784, 1985.
- 394 23. Shulman G, Corbetta M, Buckner RL, Raichle ME, Fiez JA, Miezin FM, Petersen SE. Top-down  
395 modulation of early sensory cortex. *Cerebral Cortex* 7: 193–206, 1997.
- 396 24. Veniero D, Gross J, Morand S, Duecker F, Sack AT, Thut G. Top-down control of visual cortex by  
397 the frontal eye fields through oscillatory realignment. *Nature Communications* 12, 2021.
- 398 25. Ekstrom LB, Roelfsema PR, Arsenault JT, Bonmassar G, Vanduffel W. Bottom-up dependent  
399 gating of frontal signals in early visual cortex. *Science* 321: 414-417, 2008.
- 400 26. Movshon JA, Thompson ID, Tolhurst DJ. Spatial summation in the receptive field of simple cells  
401 in the cat's striate cortex. *J Physiol* 283: 53-77, 1978.
- 402 27. Ringach DL. Mapping receptive fields in primary visual cortex. *J Physiol* 558: 717-728, 2004.
- 403 28. Nakano T, Kato M, Morito, Y, Itoi S, Kitazawa S. Blink-related momentary activation of the  
404 default mode network while viewing videos. *Proc Natl Acad Sci USA* 110: 702-706, 2013.

405

## 406 **FIGURE LEGENDS**

407 **Figure 1. Experimental design with sample stimulus frames displaying the high temporal**  
408 **contrast and low temporal contrast displays.** High temporal contrast flickered at 7 Hz with a  
409 luminance amplitude envelope of 100%, encompassing the maximum (255 a.u.) and minimum (0 a.u.)  
410 possible luminance values. The low temporal contrast events also flickered at 7 Hz with a luminance  
411 amplitude envelope of 10%, encompassing a range of luminance values between 140 a.u. and 115  
412 a.u.

413

414 **Figure 2. Eye closure has minimal effect on visual responses in LGN and V1, while suppressing**  
415 **responses in V2 and V3.** (A) Event-triggered average for luminance task across ROI and eye  
416 condition. Across LGN, V1, and V2, during eyes open runs high temporal contrast stimuli elicits a  
417 greater BOLD response than with low temporal contrast stimuli. Though there is no effect of temporal  
418 contrast in V3, BOLD response increases regardless of the stimulus temporal contrast. During eye  
419 closure, BOLD responses in LGN and V1 during the high temporal contrast stimuli elicits a similar  
420 BOLD response as during eyes open runs. With eye closure, V2 and V3 have strongly attenuated  
421 BOLD regardless of temporal contrast. Red plots indicate high temporal contrast trials and blue  
422 indicates the low temporal contrast trials. The grey bar indicates 16 second period of stimulus  
423 presentation. Error shading is 1 SEM. N=8 subjects. (B) Average BOLD activation during stimulus  
424 presentation across conditions. Pairwise comparisons show a significant decrease in V2 and V3 BOLD  
425 magnitude with eye closure for high temporal contrast stimuli. In LGN, BOLD magnitude with high  
426 temporal contrast stimuli with eyes open was marginally greater than low contrast ( $t(14)=1.79$ ;  $P =$   
427  $0.047$ ) at a Bonferroni corrected p-value cutoff of 0.0125. In V1 with eyes closed, BOLD magnitude  
428 During high temporal contrast stimuli was marginally greater than during low temporal contrast stimuli  
429 ( $t(14)=1.70$ ;  $P = 0.055$ ). In V2, BOLD magnitude with high temporal contrast stimuli with eyes open  
430 was greater than low contrast ( $t(14)=2.51$ ;  $P = 0.012$ ) and high temporal contrast stimuli with eyes  
431 closed was suppressed compared to eyes open. In V3, BOLD magnitude during high temporal contrast  
432 stimuli with eyes closed was also suppressed compared to eyes open. Y-axis is BOLD signal averaged  
433 across 4-16s post-stimulus onset. Error bars are 1 SEM. All p-values from pairwise comparison only  
434 survive multiple comparison correction at a p-value less than 0.0125, using Bonferroni correction  
435 (0.05/n where n=4 per ROI). \*  $P < 0.0125$



436

437 **Figure 3. The effects of eye closure do not depend on eccentricity tuning.** (A) Event-triggered  
438 Average for BOLD response to luminance task across cortical ROI and eye condition separated by  
439 voxels tuned to different portions of the visual field. With eyes open and eyes closed, the BOLD  
440 responses to high contrast stimuli are uniform across eccentricities for all cortical ROIs. Foveal voxels  
441 were tuned to between 0.01 dva – 1.5 dva. Parafoveal voxels were tuned to between 1.5 dva – 4.0  
442 dva. Peripheral voxels were tuned to between 4.0 dva – 17 dva. (B) Average BOLD activation during  
443 stimulus presentation across conditions (top = eyes closed; bottom = eyes open), separated by  
444 eccentricity preference. There are no significant pairwise comparisons when comparing eccentricity  
445 responses within each ROI. Y-axis is BOLD signal averaged across 4-16s post-stimulus onset. Error  
446 bars and error shading is 1 SEM.


# Eye closure produces distinct responses in early and late visual processing pathways

## Methods

Subjects viewed a temporal contrast modulated visual stimulus with eyes open or eyes closed while collecting fMRI



## Outcomes




## Conclusions

Visual information is processed in early visual processing regions with eyes open and closed, whereas eye closure attenuates visual responses in later visuocortical regions



# Sample stimulus blocks

*high temporal contrast*



*low temporal contrast*






*luminance*




Downloaded from journals.physiology.org/journal/jn at Boston Univ (128.197.062.166) on June 21, 2024.

← 16 s at a contrast modulation of 7 Hz →

← 16 s at a contrast modulation of 7 Hz →



A



B

